

# Toppers Village



## Relations and Functions

Class Notes ( pdf )

**Class 12 Maths**

YouTube पर इस Chapter के FREE Videos देखिए ...  
(Playlist की Link नीचे है) ↓↓↓

<https://www.youtube.com/watch?v=IC1Ccq7LPAE&list=PLPFrn0ppwwkR3IRWN8xWFgqbrSiyX-Btm>

Relations and Functions (संबंध एवं फलन)

Set  $A = \{a, b, c\}$  = set of cities


$a \rightarrow$  Ajmer  
 $b \rightarrow$  Bombay

Set  $B = \{r, s, t\}$  = set of states

$c \rightarrow$  Chennai  
 $r \rightarrow$  Rajasthan  
 $s \rightarrow$  Sikkim  
 $t \rightarrow$  Tamilnadu

Cartesian Product  $A \times B =$   
(कार्टीसिय गुणन)  
↓  
'set of all ordered pairs from A to B'

- $(a, r), (a, s), (a, t),$
- $(b, r), (b, s), (b, t),$
- $(c, r), (c, s), (c, t)$



Arrow Diagram

Set Builder Form

Relation ' $R_1$ ' =  $\{(x, y) : x \in A, y \in B, \text{city } x \text{ lies in state } y\}$

Relation ' $R_1$ ' =  $\{(a, r), (c, t)\}$  ← Roster Form

# Relation  $R$  is a subset of Cartesian Product  $A \times B$ :  
from A to B  $\boxed{R \subseteq A \times B}$

#  $(a, r) \in R, \forall a, r$   
 $(c, t) \in R, \forall c, t$

~~$(a, s) \in R$~~   
 $(a, s) \notin R$

# Relation A to A  
↓  
Relation on 'A'  
(in)

## Empty Relation (रिक्त संबंध) $\phi \subseteq A \times A$

$$\text{Set } A = \{1, 2, 3\}$$

$$\text{Set } B = \{100, 101, 102\}$$

$$\text{Relation } R_2 = \{(a, b) : a \in A, b \in B, \underline{a > b}\} = \{\} = \phi$$

impossible.

## Universal Relation (सार्वत्रिक संबंध) $= A \times A$

$$R_3 = \{(a, b) : a \in A, b \in B, |a - b| \geq 0\} = \left\{ \begin{array}{l} (1, 100) \\ (1, 101) \\ \vdots \\ (3, 102) \end{array} \right\}$$

$$\{1, 2, 3\}$$

$$\{100, 101, 102\}$$

100% Possible (Sure)

↑  
All

$$\begin{aligned} |1 - 100| &= |-99| \geq 0 \\ &= +99 \geq 0 \quad \text{(True)} \end{aligned}$$

## Different Types of relations on 'A'


Relation 'R' from A to A  $\Rightarrow$  R on 'A'

Extra

I Identity Relation:

$$A = \{1, 2, 3\} \longrightarrow A = \{1, 2, 3\}$$

$$I_A = \{(1, 1), (2, 2), (3, 3)\} \checkmark$$



$$\times R_1 = \{(1, 1), (2, 2)\}$$

$$\times R_2 = \{(1, 1), (2, 2), (3, 3), (1, 2)\}$$

II

Reflexive Relations

on A

A to A  
{1,2,3} to {1,2,3}

if (a,a) ∈ R ∀ a ∈ A  
(for every)

A = {1,2,3}

R1 = {(1,1), (2,2), (3,3)} Reflexive ✓  
Identity ✓

R2 = {(1,1), (2,2)} Reflexive X  
Identity X

R3 = {(1,1), (2,2), (3,3), (1,2)} Reflexive ✓  
Identity X

R4 = {(1,1), (2,2), (3,3), (1,2), (2,3), (3,1)} Reflexive ✓  
Id. X

III

Symmetric Relations (सममित संबंध)

A to A  
{1,2,3} to {1,2,3} R ⇒ if (a,b) ∈ R then (b,a) ∈ R

R1 = {(1,1), (2,2)} ✓ (1,1) ∈ R1, (1,1) ∈ R1

R2 = {(1,1), (2,2), (3,3)} ✓

R3 = {(1,1), (2,2), (3,3), (1,2)} X

R4 = {(1,2), (2,1), (3,2)} Sym. X

R5 = {(1,2), (2,1)} Sym. ✓

## IV Transitive Relations (संक्रामक संबंध)

if  $(a,b) \in R, (b,c) \in R$  then  $(a,c) \in R$

Transitive  $\times$

$$A = \{1, 2, 3\} \longrightarrow A = \{1, 2, 3\}$$

$$R_1 = \{(1,2), (2,3), (1,3)\} \quad (1,2) \in R_1, (2,3) \in R_1 \Rightarrow (1,3) \in R_1$$

$$R_2 = \{(1,1)\} \quad (1,1) \in R_2, (1,1) \in R_2 \Rightarrow (1,1) \in R_2$$

$$R_3 = \{(1,1), (2,2), (3,3)\}$$

$$R_4 = \{(1,2)\}$$

If rain then play

$$R_5 = \{(1,1), (2,2), (3,3), (1,2)\}$$

$$\times R_6 = \{(1,2), (2,3), (3,1)\}$$

$$(1,3) \notin R_6$$

## V Equivalence Relation (समतुल्य संबंध)

Reflexive  $(a,a) \in R \forall a \in A$

Symmetric  $(a,b) \Rightarrow (b,a)$

Transitive  $(a,b), (b,c) \Rightarrow (a,c)$

e.g.

$$A = \{1, 2, 3\}$$

$$R : A \rightarrow A$$

$$R = \{(1,1), (2,2), (3,3), (1,2), (2,1), (2,3), (3,2), (1,3), (3,1)\}$$

(I) Reflexive  $(a,a) \in R \quad \forall a \in A$  ✓

(II) Symmetric ✓

$$(a,b) \Rightarrow (b,a)$$

(III) Transitive ✓

If  $(a,b) \in R, (b,c) \in R$   
then  $(a,c) \in R$

Equivalence  
Relation

Exercise 1.1      [Relations and Functions]

$R$  ~~for~~ from 'A' to 'B'

Reflexive Relations      in 'A' on 'A' = from A to A


$(a,a) \in R$  for every  $a \in A$ .

Symmetric Relations

if  $(a,b) \in R$  then  $(b,a) \in R$ .

Transitive Relations

If  $(a,b) \in R$  and  $(b,c) \in R$  then  $(a,c) \in R$ .



Equivalence Relation

Exercise 1.1

[Q.1] Reflexive / Symmetric / Transitive

(i)  $A = \{1, 2, 3, \dots, 13, 14\}$  (in A)  $x \in A, y \in A$

$R = \{(x,y) : 3x - y = 0\}$

Reflexive for  $(x,x)$        $3x - x = 0 \Rightarrow 2x = 0 \Rightarrow \boxed{x=0} \notin A$

Not reflexive,       $(0,0)$        $x \notin A$

Symmetric

If  $(x,y) \in R$  then  $(y,x) \notin R$

Not Sym.

$3x - y = 0$

$\nRightarrow$

$3y - x = 0$

Transitive,

If  $(x,y) \in R$  and  $(y,z) \in R$  then  $(x,z) \in R$

$$\begin{array}{ccc} \downarrow & & \downarrow \\ 3x - y = 0 & & 3y - z = 0 \\ \textcircled{1} & & \textcircled{2} \end{array} \quad \left. \vphantom{\begin{array}{ccc} \downarrow & & \downarrow \\ 3x - y = 0 & & 3y - z = 0 \\ \textcircled{1} & & \textcircled{2} \end{array}} \right\} \boxed{3x - z = 0} ??$$

$3x = y \Rightarrow 3(3x) - z = 0$   
 $\boxed{9x - z = 0}$

Not Transitive

(ii) Relation in  $\mathbb{N}$  (set of natural no.)  
(from  $\mathbb{N}$  to  $\mathbb{N}$ )  $\{1, 2, 3, \dots\}$

$$R = \{ (x, y) : y = x + 5 \text{ and } x < 4 \}$$

$$x = 1, 2, 3 \quad y = 6, 7, 8$$

Set Builder Form

$$R = \{ (1, 6), (2, 7), (3, 8) \}$$

Reflexive

$$\textcircled{(1, 1) (2, 2) (3, 3) (4, 4) \dots}$$

Not reflexive

Symmetric

$(1, 6) \in R$  but  $(6, 1) \notin R$   
 $\therefore$  Not sym.

Transitive,

$$\begin{array}{l} (1, 6) \in R \\ (2, 7) \in R \\ (3, 8) \in R \\ \text{only} \end{array}$$

Transitive,



(iii) Relation in the set  $A = \{1, 2, 3, 4, 5, 6\}$

$$R = \{(x, y) : \underline{y \text{ is divisible by } x}\}$$

Reflexive.

$$\cancel{(x, x)} \in R \quad \forall x \in A$$

↓

$x$  is divisible by  $x \quad \forall x \in A$

(True)

Yes: Reflexive ✓

Symmetric: If  $(x, y) \in R$  then  $(y, x) \in R$ .

$y$  is divisible by  $x$

$$6 \text{ --- } || \text{ --- } 2$$

✓

$x$  is divisible by  $y$

$$2 \text{ --- } || \text{ --- } 6$$

X

(Not Sym.)

Transitive. If  $(x, y) \in R$  and  $(y, z) \in R$  then  $(x, z) \in R$

$y$  is divisible by  $x$

$z$  is divisible by  $y$

$z$  is divi.  
by  $x$

$$y = kx$$

$$z = m \cdot y$$

$$\Rightarrow z = \underline{mk} \cdot x$$

$$z = \underline{mk} \cdot x$$

Yes Transitive.



(v) relation  $R$  in set 'A' of human beings in a town.

(a)  $R = \{(x, y) : x \text{ and } y \text{ work at the same place}\}$

(b)  $R = \{(x, y) : x \text{ and } y \text{ live in the same locality}\}$

Reflexive.

$$(x, x) \in R$$

✓

Symmetric

$$(x, y) \in R \Rightarrow (y, x) \in R$$

✓

✓

Transitive.

$$(x, y) \in R, (y, z) \in R$$

$\Rightarrow$

$$(x, z) \in R$$

✓

(c)  $R = \{(x, y) : x \text{ is exactly } 7\text{cm taller than } y\}$

Reflexive

$$(x, x) \notin R$$

Not

✓

Sym

$$(x, y) \in R$$

$\Rightarrow$

$$(y, x) \notin R$$

X

Not

✓

Transitive.

$$(x, y) \in R \text{ \& } (y, z) \in R$$

$\Rightarrow$

$$(x, z) \notin R$$

$$x = z + 7$$

$$x = y + 7$$

$$x = z + 7$$

$$y = z + 7$$

$$x = (z + 7) + 7$$

$$x = z + 14$$

Not

(d)  $R = \{(x, y) : x \text{ is wife of } y\}$

Reflexive


$$(x, x) \in R$$

X

Not

Sym.

$$(x, y) \in R \Rightarrow (y, x) \in R$$



Transitive

$$(x, y) \in R, (y, z) \notin R \Rightarrow (x, z) \in R$$

Transitive

$$\textcircled{c} R = \{(x, y) : x \text{ is father of } y\}$$

Reflexive

$$(x, x) \in R$$

X

Sym.

$$(x, y) \in R \Rightarrow (y, x) \in R$$

X

Transitive

$$(x, y) \in R, (y, z) \in R \Rightarrow (x, z) \notin R$$



Grand Father

Father

X

toppers village

Exercise 1.1

Relations and Functions


Q.2 Relation in the set 'R' (real numbers)

$R = \{(a,b) : a \leq b^2\}$  } Reflexive ✗  
 } Symmetric ✗  
 } Transitive ✗ } Prove

Reflexive

$(x,x) \in R \quad \forall x \in R$

relation  
for every




$\Rightarrow x \leq x^2$

$x = 0.5 = \frac{1}{2}$

$\frac{1}{2} \neq (\frac{1}{2})^2$

$\frac{1}{2} \neq \frac{1}{4}$

$0.5 > 0.25$



this is not true  
for  $x \in (0,1)$

Not reflexive

Symmetric

If  $(x,y) \in R$  then  $(y,x) \in R$

$x \leq y^2$

$y \leq x^2$

$x=1, y=2$

$1 \leq 4$

Not Sym.

$2 \leq 1$

✗

Transitive

If  $(a,b) \in R, (b,c) \in R$  then  $(a,c) \in R$

$a \leq b^2$

$b \leq c^2$

$a \leq c^2$

Not

$7 \leq 25$

$a=7, b=-5$

$c=1$

$7 \leq 1$  ✗

$(-5 \leq 1)$

Q.3 Relation (R) in the set  $A = \{1, 2, 3, 4, 5, 6\}$

$$R = \{(a, b) : b = a + 1\}$$



Reflexive

$$(a, a) \in R \quad \forall a \in A$$

$$a = a + 1$$

$$\Rightarrow 0 = 1$$

False

Not Reflexive

Symmetric

$$(a, b) \in R \Rightarrow (b, a) \in R$$

$$b = a + 1$$

$$a = b + 1$$



X

Not

Transitive


$$(a, b) \in R, (b, c) \in R$$

$$\Rightarrow (a, c) \in R$$

$$b = a + 1 \quad \text{--- (1)}$$

$$c = b + 1 \quad \text{--- (2)}$$

$$+ \quad c = a + 2$$



Q.4 Relation 'R' in set 'R' ← Real no.

$$R = \{(a, b) : a \leq b\}$$

Reflexive ✓  
Transitive ✓  
Sym. X

Reflexive

$$(a, a) \in R$$

$$a \leq a$$

✓  
✓

Sym.

$$(a, b) \in R \Rightarrow (b, a) \in R$$

$$a \leq b$$

$$b \leq a$$

$$\downarrow \quad \downarrow$$

$$2 \leq 5$$

$$5 \leq 2$$


Not Sym.

Transitive

$$(a, b) \in R, (b, c) \in R$$

$$a \leq b$$

$$b \leq c$$



$$(a, c) \in R$$

$$a \leq c$$

Transitive

Q.5 relation R in  $\mathbb{R} \rightarrow$  real no.

$$R = \{(a,b) : a \leq b^3\}$$



Reflexive

$$(a,a) \in R \quad \forall a \in \mathbb{R}$$

$$a \neq a^3$$

$$a = \frac{1}{2}$$

$$\frac{1}{2} \neq \left(\frac{1}{2}\right)^3$$

$$\frac{1}{2} \neq \frac{1}{8}$$

$$0.5 > 0.125$$

Not

Transitive

$$(a,b) \in R, (b,c) \in R$$

$$a \leq b^3 \quad b \leq c^3$$

$$(a,c) \notin R$$

$$a \neq c^3$$

$$a \leq (b)^3 \leq (c^3)^3$$

$$a \leq c^9$$

Not transitive

Symmetric

$$(a,b) \in R \Rightarrow (b,a) \in R$$

$$a \leq b^3$$

$$b \leq a^3$$

$$a = 1$$

$$b = 2$$

Not Sym.

Q.6 Relation R in the set  $\{1,2,3\}$

$$R = \{(1,2), (2,1)\}$$

Sym. ✓

Reflexive ✗

Trans. ✗

Reflexive

$$\{(1,1), (2,2), (3,3)\}$$

Sym.

$$(1,2) \in R \text{ then } (2,1) \in R$$

Transitive,  $(a,b), (b,c) \Rightarrow (a,c)$

$$(1,2), (2,1) \Rightarrow (1,1) \notin R$$

Not transitive

in set 'A' of all books in a library.

Q.7

$$R = \left\{ (x, y) : x \text{ and } y \text{ have same no. of pages} \right\}$$

Equivalence

$$(x, x) \rightarrow \checkmark$$

$$\left[ \begin{array}{l} \bullet \text{ Reflexive} \\ \bullet \text{ Symmetric} \\ \bullet \text{ Transitive} \end{array} \right] \rightarrow (x, y) \in R \Rightarrow (y, x) \in R \quad \checkmark$$

$$\rightarrow (x, y) \in R \text{ and } (y, z) \in R \Rightarrow (x, z) \in R \quad \checkmark$$


Q.8

Relation in set  $A = \{1, 2, 3, 4, 5\}$

$$R = \left\{ (a, b) : |a - b| \text{ is even} \right\}$$

$$R = \left\{ \begin{array}{l} (1, 1), (1, 3), (1, 5) \\ (2, 2), (2, 4) \\ (3, 1), (3, 3), (3, 5) \\ (4, 2), (4, 4) \\ (5, 1), (5, 3), (5, 5) \end{array} \right\}$$

$\{1, 3, 5\}$



$$\begin{array}{l} (\text{odd} - \text{odd}) = \text{even} \\ | \text{even} - \text{even} | = \text{even} \end{array}$$

Reflexive

$$(a, a) \in R \quad \forall a \in A$$

Symmetric

$$(a, b) \Rightarrow (b, a)$$

Transitive

$$(a, b), (b, c) \Rightarrow (a, c)$$



Q.9 Relation in the set  $A = \{x \in \mathbb{Z} : 0 \leq x \leq 12\}$   
 $A = \{0, 1, 2, 3, \dots, 12\}$

(i)  $R = \{(a, b) : |a-b| \text{ is a multiple of } 4\}$

(ii)  $R = \{(a, b) : a = b\}$

(i)  $R = \{(a, b) : |a-b| \text{ is a multiple of } 4\}$

Reflexive.  $(a, a) \in R \quad \forall a \in A$

$\Rightarrow |a-a| = 0$  is a multiple of 4

Sym.  $(a, b) \in R \Rightarrow (b, a) \in R$

$|a-b|$  is a multiple of 4

$|b-a|$  is a multiple of 4

True.

Transitive.  $(a, b) \in R, (b, c) \in R \Rightarrow (a, c) \in R$

$|a-b| = 4k \quad |b-c| = 4m \quad \Rightarrow \quad |a-c| = 4n$

$a-b = 4k \quad b-c = 4m$


$\rightarrow (+) \checkmark$

$a-c = 4k + 4m$

$a-c = 4(k+m)$

$|a-c| = 4(k+m)$

elements related to 1



c)  $(a-b)$  is a multiple of 4

0, 4, 8, 12, ...

$|1-b|$  is a multiple of 4


$b = 1, 5, 9$

cii)  $a=b$

$(a, a)$  —

$(a, b) (b, a)$  —

$(a, b) (b, c) (a, c)$  —



Toppers Village Class 12 Maths

Exercise 1.1 (Relations and Functions)

Q11 to Q16

Q.11 Relation in the set A of points in a plane.

$R = \{ (P, Q) : \text{distance of the point } P \text{ from origin} \}$   
 $\downarrow$  is same as the distance of the  
 $OP = OQ$  Point  $Q$  from origin  $\}$

Reflexive

$(P, P) \in R \quad \forall P \in A$

$\downarrow$   
 origin = 'O' (0,0)

$OP = OP$   
 ✓ True.

Reflexive  
 ✓  
 ✓

Symmetric

$(P, Q) \in R \Rightarrow (Q, P) \in R$

$\downarrow \quad \quad \quad \uparrow$   
 $OP = OQ \quad OQ = OP$

✓  
 ✓

Transitive


$(P, Q) \in R, (Q, S) \in R$

$OP = OQ \quad \quad \quad OQ = OS$   
 ①  $\quad \quad \quad$  ②


$\Downarrow$   
 $(P, S) \in R$   
 $OP = OS$

✓  
 ✓

Equivalence



Sides



Q.12 Relation in the set A of all triangles.

$R = \{ (T_1, T_2) : T_1 \text{ is similar to } T_2 \} \rightarrow T_1 \sim T_2$

Reflexive

$T_1 \sim T_1$   
 ✓

Symmetric

$T_1 \sim T_2 \Rightarrow T_2 \sim T_1$   
 ✓

Transitive

$T_1 \sim T_2, T_2 \sim T_3 \Rightarrow T_1 \sim T_3$   
 ✓

**Q.13** Relation in the set  $A$  of all polygons

$$R = \{(P_1, P_2) : P_1 \text{ and } P_2 \text{ have same no. of sides}\}$$

(let:  $n(P)$  denotes no. of sides of 'P')

Reflexive.  
 $(P_1, P_1) \in R \forall P_1 \in A$   
 $n(P_1) = n(P_1)$   
 ✓

Sym.  
 $(P_1, P_2) \in R \Rightarrow (P_2, P_1) \in R$   
 $n(P_1) = n(P_2) \quad n(P_2) = n(P_1)$   
 ✓

Transitive  $(P_1, P_2) \in R, (P_2, P_3) \in R \Rightarrow (P_1, P_3) \in R$   
 $n(P_1) = n(P_2)$  (1),  $n(P_2) = n(P_3)$  (2)  $\therefore n(P_1) = n(P_3)$   
 ✓

**Q.14** Relation in set L of all lines in XY plane.

$$R = \{(L_1, L_2) : L_1 \text{ is parallel to } L_2\}$$

Reflexive  
 $(L_1, L_1) \in R$   
 $L_1 \parallel L_1$   
 ✓

Sym.  
 $(L_1, L_2) \in R \Rightarrow (L_2, L_1) \in R$   
 $L_1 \parallel L_2 \quad L_2 \parallel L_1$   
 ✓

Transitive  
 $(L_1, L_2), (L_2, L_3) \Rightarrow (L_1, L_3)$   
 $L_1 \parallel L_2 \quad L_2 \parallel L_3 \Rightarrow L_1 \parallel L_3$   
 $\begin{matrix} \xrightarrow{m} L_1 \\ \xrightarrow{m} L_2 \\ \xrightarrow{m} L_3 \end{matrix}$   
 $y = mx + c$   
 Slope

Equivalence ✓

$$y = 2x + 4 \parallel y = 2x + c \quad c \in R$$

Q.15

Relation in the set  $\{1, 2, 3, 4\} = A$

$$R = \{(1, 2), (2, 2), (1, 1), (4, 4), (1, 3), (3, 3), (3, 2)\}$$

Reflexive

$(a, a) \in R$   
 $\forall a \in A$   
✓

Symmetric

Here  
 $(1, 2) \in R$   
but  $(2, 1) \notin R$

Transitive

$(1, 2) \in R$     $(2, 2) \in R$   
 $(1, 2) \in R$  ✓

$(a, b), (b, c) \Rightarrow (a, c)$  ✓

↓  
Option (B)

Q.16

Relation in the set N ✓

$$R = \{(a, b) : a = b - 2, b > 6\}$$

(A)  $(2, 4) \in R$   
✗  $4 < 6$

(B)  $(3, 8) \in R$   
 $3 \neq 8 - 2$

(C)  $(6, 8) \in R$   
 $6 = 8 - 2$   
✓

(D)  $(8, 7) \in R$   
 $8 \neq 7 - 2$

Option (C)

- One-one Function (Injective Fn.) (एकैकी फलन)
- many-one Function (अहुएकैकी फलन)
- Onto Function (Surjective Fn.) (आच्छादक फलन)
- Into Function (अ-आच्छादक फलन)
- ⇒ one one onto Fn. (Bijective Fn.)

Function (फलन)

Function is a special type of relation from a non empty set A to another non empty set B such that each element of set A has unique image in set B.








image of '1' = a  
Preimage of 'a' = 1

function  $f: A \rightarrow B$

Domain  
input (x)

Codomain

Range = set of images  
output (y = f(x))

Range  $\subseteq$  Codomain

# One One Functions (Injective Fn.) (एकक प्रफल)

$$f: X \rightarrow Y$$

$$0 \rightarrow 0$$


images of distinct elements of  $X$  under ' $f$ ' are distinct

$$x_1, x_2 \in X$$


$$f(x_1) = f(x_2) \Rightarrow x_1 = x_2 \quad \text{only}$$

$$x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2)$$


Many-one Functions : which are not one-one.



one-one




many-one





one-one

Horizontal line Test : (for graphs)


If a Horizontal line cuts the graph of a function at atmost  $(\sqrt{y_1}, \sqrt{y_1})$  one point, then this graph represents a one-one function.



many-one



one-one



many-one Fn.

e.g. Check injectivity (one-one / many-one)

(i)  $f: \mathbb{R} \rightarrow \mathbb{R}$

$f(x) = 3x + 7$

Let  $x_1, x_2 \in \mathbb{R}$

$f(x_1) = f(x_2)$

$\Rightarrow 3x_1 + 7 = 3x_2 + 7$

$\Rightarrow \boxed{x_1 = x_2}$  only.

one-one function

(ii)  $f: \mathbb{R} \rightarrow \mathbb{R}$

$f(x) = x^2$

Let  $x_1, x_2 \in \mathbb{R}$

$f(x_1) = f(x_2)$

$\Rightarrow x_1^2 = x_2^2$


$\Rightarrow \boxed{x_1 = \pm x_2}$  (many-one)

$\swarrow$   
 $\boxed{x_1 = x_2}$





$\searrow$   
 $\boxed{x_1 = -x_2}$

Onto Functions (Surjective Function) (आच्छादक फलन)  
(पर)  $f: X \rightarrow Y$

if every element of  $Y$ , is the image of some element of  $X$  under  $f$ .




(अं) Into Functions : which are not onto.






# Bijjective Functions (one-one onto Functions)

which are one one & onto both.



e.g. Check Surjectivity (Hence Bijjectivity)

onto/into




$x \in \mathbb{R} \leftarrow \text{Domain}$

$3x + 7 \in \mathbb{R}$

$f(x) \in \mathbb{R} \leftarrow \text{Range}$

output  $\text{Range} = \mathbb{R} = \text{Codomain}$

ONTO  
 one-one Bijjective




Range

$x \in \mathbb{R}$

$x^2 \geq 0$

Square  $\geq 0$

$x^2 \in [0, \infty)$



$f(x) \in [0, \infty) \leftarrow \text{Range}$   
 output

$\text{Range} = [0, \infty) \neq \mathbb{R}$   
 (-∞, ∞)

into many-one

Tip \* Questions with  $f: \mathbb{N} \rightarrow \mathbb{N}$

Draw Arrow Diagram

1, 2, 3, ...

\* Questions with  $f: \mathbb{R} \rightarrow \mathbb{R}$

Apply Proper methods

$f(x_1) = f(x_2)$


(one-one)  $\rightarrow x_1 = x_2$

Range = Codomain

(ONTO)

Exercise 1.2 [Relations and Functions]

Q.1  $f: \overset{\text{Domain}}{\mathbb{R}^*} \rightarrow \overset{\text{Codomain}}{\mathbb{R}^*}$  defined by  $f(x) = \frac{1}{x}$



One-one  $\stackrel{\text{let}}{=} f(x_1) = f(x_2) \xrightarrow{\text{Prove}} \boxed{x_1 = x_2} \stackrel{\text{only}}{=}$

$\Rightarrow \frac{1}{x_1} = \frac{1}{x_2}$

$\Rightarrow x_1 = x_2 \text{ only}$


$\therefore f \rightarrow \text{one-one}$  ✓

onto Range = codomain ( $\mathbb{R}^*$ ) Prove

really Output

$x \rightarrow$  input

$\exists, f(x) \rightarrow$  output = Range



onto proved



$\frac{1}{0} = \infty$   
 $\frac{1}{\infty} = 0$

Range

$f(x) = \frac{1}{x}$

Domain (input)

$x \in (-\infty, 0) \cup (0, \infty)$

$\frac{1}{x} \in (-\infty, 0) \cup (0, \infty)$

$f(x) \in (-\infty, 0) \cup (0, \infty)$


Output

Range =  $(-\infty, 0) \cup (0, \infty)$   
 =  $(-\infty, \infty) - \{0\}$   
 =  $\mathbb{R} - \{0\}$   
 =  $\mathbb{R}^* = \text{Codomain}$

Original function  $f: \mathbb{R}_* \rightarrow \mathbb{R}_*$ ,  $f(x) = \frac{1}{x}$  one-one onto

Next Case  $f: \mathbb{N} \rightarrow \mathbb{R}_*$ ,  $f(x) = \frac{1}{x}$   $\begin{matrix} \rightarrow ? \\ \rightarrow ? \end{matrix}$   
 $\{1, 2, 3, \dots\}$

Arrow Diagram.



$$f(1) = \frac{1}{1} = 1$$

$$f(2) = \frac{1}{2} = 0.5$$

$$f(3) = \frac{1}{3} = 0.\bar{3}$$

$$f(4) = \frac{1}{4} = 0.25$$

one-one  
✓

onto  $\rightarrow$  X  
 $\mathbb{R}_*$  has all elements


connected

into

Range  $\neq$  Codomain  
 $\mathbb{R}_*$

**Q.2** Check injectivity & Surjectivity  
 (one-one) (onto)  
 (manyone) (into)

(i)  $f: \mathbb{N} \rightarrow \mathbb{N}$  given by  $f(x) = x^2$




$f(1) = 1^2 = 1$   
 $f(2) = 2^2 = 4$

one-one ✓ | onto X  
 Range  $\neq$  Codomain  
 $\downarrow$   $\downarrow$   
 $\mathbb{N}$   
 $\{1, 2, 3, \dots\}$

one-one & into

(ii)  $f: \mathbb{Z} \rightarrow \mathbb{Z}$  given by  $f(x) = x^2$

↓  
 integers



$f(0) = 0^2 = 0$   
 $f(-1) = (-1)^2 = 1$   
 $f(-2) = (-2)^2 = 4$   
 $f(1) = 1^2 = 1$   
 $f(2) = 2^2 = 4$

Surjectivity

Range  $\neq$  Codomain  
 $\downarrow$

$\{0, 1, 4, 9, 16, \dots\} \neq \mathbb{Z}$

~~one~~ Injectivity

many-one ✓

into

(iii)  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) = x^2$   
 $(-\infty, \infty)$

One-one  $x_1, x_2 \in \mathbb{R}$  ??

$f(x_1) = f(x_2) \rightarrow x_1 = x_2$  only

$\Rightarrow x_1^2 = x_2^2$

$\Rightarrow x_1 = \pm x_2$

$x_1 = x_2$  or  $x_1 = -x_2$

many one

onto Range = Codomain  $\mathbb{R}$

output

$f(x)$

$f(x) = x^2$

$x \in \mathbb{R}$

$x^2 \geq 0$

$f(x) \geq 0$

$f(x) \in [0, \infty)$


Range =  $[0, \infty)$

Codomain =  $\mathbb{R} = (-\infty, \infty)$

Range  $\neq$  Codomain

Into

(iv)  $f: \mathbb{N} \rightarrow \mathbb{N}$  given by  $f(x) = x^3$



one-one



into

$f(x) = x^3$

$f(1) = 1^3 = 1$

$f(2) = 2^3 = 8$


Surjectivity

Range  $\neq$  Codomain

$\{1, 8, 27, 64, \dots\} \neq \{1, 2, 3, \dots\}$

⑤  $f: \mathbb{Z} \rightarrow \mathbb{Z}$  given by  $f(x) = x^3$

$\{-2, -1, 0, 1, 2, 3, \dots\}$



$$f(x) = x^3$$

$$f(0) = 0$$

$$f(1) = 1$$

$$f(2) = 8$$

$$f(3) = 27$$

$$f(-1) = -1$$

$$f(-2) = -8$$

$$f(-3) = -27$$

one-one  
✓

onto  
✗

into  
✓

Range  $\neq$  Codomain

$\{\dots, 27, 8, 1, 0, -1, -8, \dots\} \neq \mathbb{Z}$

Toppers Village Class 12 maths

Exercise 1.2 (Relations and Functions)

**Q.3** Prove that the greatest integer function  $f: \mathbb{R} \rightarrow \mathbb{R}$ , given by  $f(x) = [x]$ , is neither one-one nor onto.

one-one  $f: \mathbb{R} \rightarrow \mathbb{R}$   
 $f(x) = [x]$   $\uparrow$   
 $x$   
input

$x_1 = 8.2$   
 $f(x_1) = f(8.2) = [8.2] = 8$


$x_2 = 8.5$   $\parallel$   
 $f(x_2) = f(8.5) = [8.5] = 8$

$x_1 \neq x_2$   
but  $f(x_1) = f(x_2)$


not one-one

Greatest integer function  
 $f(x) = [x]$


$f(2) = [2] = 2$   
 $f(2.3) = [2.3] = 2$



$f(0.02) = [0.02] = 0$



$f(-1.7) = [-1.7] = -2$



onto / into  
 $\downarrow$

Range = Codomain  
 $\downarrow$   $\mathbb{R}$

actual output

$f(x) = [x] = \text{Integer} = \mathbb{Z}$

Range =  $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$

Range =  $\mathbb{Z}$   $\times$   
Codomain =  $\mathbb{R}$

into

Q.4  $f: \mathbb{R} \rightarrow \mathbb{R}$   $f(x) = |x|$

neither one-one nor onto

one-one.



$f(x) = f(x_2) \Rightarrow$   $x_1 = x_2$  only??

$\Rightarrow |x_1| = |x_2|$

$\Rightarrow x_1 = \pm x_2$

$x_1 = x_2$   
 $x_1 = -x_2$

Not one-one




onto

Range = Codomain  $(\mathbb{R})$

output

$f(x) = |x|$

$|x| \geq 0$



Range =  $[0, \infty)$

Codomain =  $\mathbb{R} = (-\infty, \infty)$

Into

not onto


Q.5  $f: \mathbb{R} \rightarrow \mathbb{R}$   $f(x) = \begin{cases} 1 & x > 0 \\ 0 & x = 0 \\ -1 & x < 0 \end{cases}$  (Signum)

neither one-one nor onto

Not one-one

$f(2) = 1, f(3) = 1$

$f(-2) = -1, f(-3) = -1$



not onto

Range  $\neq$  Codomain  $(\mathbb{R})$

output

$\{1, 0, -1\}$

not onto




Q.6.

$$A = \{1, 2, 3\}$$

$$B = \{4, 5, 6, 7\}$$

$$f = \{(1, 4), (2, 5), (3, 6)\} \quad f: \underline{A} \rightarrow \underline{B}$$

Show that  $f$  is one-one



Q.7 (i)  $f: \underline{R} \rightarrow \underline{R}$  defined by  $f(x) = 3 - 4x$

one-one,

$$\text{let } f(x_1) = f(x_2) \Rightarrow x_1 = x_2 \text{ only}$$

$$f(x_1) = f(x_2)$$

$$\Rightarrow 3 - 4x_1 = 3 - 4x_2$$

$$\Rightarrow \boxed{x_1 = x_2} \text{ only}$$

One-one

Bijjective

onto

Range = Codomain

Output  
( $f(x)$ )

$$x \in R$$

$$\underline{3 - 4x} \in R$$

$$f(x) \in R$$

$$\underline{\text{Range} = R = \underline{\text{Codomain}}}$$

onto

(ii)  $f: \mathbb{R} \rightarrow \mathbb{R}$  defined by  $f(x) = 1+x^2$

$$\underline{f(x) = 1+x^2}$$

one-one

$$\text{let } f(x_1) = f(x_2)$$

$$\Rightarrow 1+x_1^2 = 1+x_2^2$$

$$\Rightarrow x_1^2 = x_2^2$$

$$\Rightarrow \boxed{x_1 = \pm x_2}$$

$$\underline{(x_1 = x_2)} \text{ या } \underline{x_1 = -x_2}$$


not one-one

not ~~Bij~~  
Bijection

onto

Range = Codomain

??



$$x \in \mathbb{R}$$

$$x^2 \geq 0$$

$$\underline{1+x^2} \geq \underline{1+0}$$

$$f(x) \geq 1$$

$-\infty \quad 1 \quad \infty$

$$f(x) \in [1, \infty)$$

Range

Range  $\neq$  Codomain

not onto

Toppers Village class 12 maths

Exercise 1.2 (Relations and Functions)


**Q.8** Let A and B be sets. show that  $f: A \times B \rightarrow (B \times A)$  such that  $f(a, b) = (b, a)$  is bijjective function. one-one & onto

Ans. Let  $A = \{a_1, a_2\}$      $B = \{b_1, b_2\}$

$A \times B = \{ \underline{(a_1, b_1)}, \underline{(a_1, b_2)}, \underline{(a_2, b_1)}, \underline{(a_2, b_2)} \}$   
 Cartesian Product


$B \times A = \{ \underline{(b_1, a_1)}, \underline{(b_2, a_1)}, \underline{(b_1, a_2)}, \underline{(b_2, a_2)} \}$

Arrow Diagram



$f(a, b) = (b, a)$   
 $f(x, y) = (y, x)$   
 $f(1, -3) = (-3, 1)$   
 $f(\underline{a_1, b_1}) = \underline{(b_1, a_1)}$

One-one



onto


$\text{Range} = \text{Codomain}$   
 $\downarrow$                        $\downarrow$   
 $B \times A = B \times A$

Bijjective  $F_n^m$

Q.9

$f: \mathbb{N} \rightarrow \mathbb{N}$  be defined by

$$f(n) = \begin{cases} \frac{n+1}{2}, & \text{if } n \text{ is odd} \\ \frac{n}{2}, & \text{if } n \text{ is even} \end{cases}; n \in \mathbb{N}$$



$$f(1) = \frac{1+1}{2} = 1$$

$$f(2) = \frac{2}{2} = 1$$

$$f(3) = \frac{3+1}{2} = 2$$

$$f(4) = \frac{4}{2} = 2$$

One-one  $\rightarrow$  Not

(many-one) ✓

$$n_1 = 1, n_2 = 2$$

$$n_1 \neq n_2 \leftarrow$$

$$\left[ \begin{array}{l} f(n_1) = f(n_2) \\ f(1) = f(2) \\ \downarrow \quad \downarrow \\ 1 \quad 1 \end{array} \right]$$

Onto / Into

Range = Codomain

$$\text{Output} \\ \{1, 2, 3, \dots\} \\ \Downarrow \\ \mathbb{N} \checkmark$$

Onto ✓

not one one

onto

Not Bijective

Q.10

$R = \{3\}$

$R = \{1\}$

$f: A \rightarrow B$  defined by  $f(x) = \frac{x-2}{x-3}$

one-one

Let  $f(x_1) = f(x_2) \Rightarrow \boxed{x_1 = x_2}$  ??

$\downarrow$   
Let  $f(x_1) = f(x_2)$

$$\Rightarrow \frac{x_1 - 2}{x_1 - 3} = \frac{x_2 - 2}{x_2 - 3}$$

$$\Rightarrow (x_1 - 2) \cdot (x_2 - 3) = (x_1 - 3) \cdot (x_2 - 2)$$

$$\Rightarrow \cancel{x_1 x_2} - 3x_1 - 2x_2 + 6 = \cancel{x_1 x_2} - 2x_1 - 3x_2 + 6$$

$$\Rightarrow -3x_1 + 2x_1 = 2x_2 - 3x_2$$

$$\Rightarrow -x_1 = -x_2 \Rightarrow \boxed{x_1 = x_2} \text{ only}$$

$\therefore f(x) \rightarrow$  one-one

onto

Range = Codomain

output

$f(x) = y$

$B = R - \{1\}$

$$y = f(x) = \frac{x-2}{x-3}$$

$x-3 \neq 0$   
 $x \neq 3$

represent 'x' in terms of 'y'

$$\Rightarrow xy - 3y = x - 2$$

$$\Rightarrow \underline{xy - x} = 3y - 2$$

$$x(y-1) = 3y - 2$$

$$\Rightarrow x = \frac{3y-2}{y-1} \neq 0$$

$$y-1 \neq 0$$

$$y \neq 1 \quad y \in R - \{1\}$$

$$\text{Range} = R - \{1\}$$

$$\text{Codomain} = R - \{1\}$$

$$\text{Range} = \text{codomain}$$

$f(x) \rightarrow$  onto



Composite Function & Invertible Function

(संयुक्त फलन)

(प्रतिलोमीय फलन)


Note: Identity Function (तत्समक फलन) (I)

$f(x) = x$   
 $y = x$


$f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) = x$  ( $= I_{\mathbb{R}}$ )

$f: \mathbb{N} \rightarrow \mathbb{N}$  given by  $f(x) = x$  ( $= I_{\mathbb{N}}$ )

$f: A \rightarrow A$  given by  $f(x) = x$  ( $= I_A$ )




Composite Function (संयुक्त फलन)  $\Rightarrow$



Definition: Let  $f: A \rightarrow B$  and  $g: B \rightarrow C$  be two functions. Then composition of 'f' and 'g' denoted by  $g \circ f$ , is defined as the function  $g \circ f: A \rightarrow C$  given by

$g \circ f(x) = g(f(x)), \quad \forall x \in A$

e.g. Let  $f: \overset{A}{\{2, 3, 4, 5\}} \rightarrow \overset{B}{\{3, 4, 5, 9\}}$  and  $g: \overset{B}{\{3, 4, 5, 9\}} \rightarrow \overset{C}{\{7, 11, 15\}}$  be defined as  $f = \{(2, 3), (3, 4), (4, 5), (5, 5)\}$  and  $g = \{(3, 7), (4, 7), (5, 11), (9, 11)\}$ . Find  $g \circ f$ .



$$\begin{array}{c} B \rightarrow C \\ \uparrow \\ A \rightarrow B \\ \uparrow \\ \underline{g \circ f} = \{(2, 7), (3, 7), (4, 11), (5, 11)\} \end{array}$$

$$\begin{aligned} (2, 3) \in f &\Rightarrow f(2) = 3 \\ (3, 7) \in g &\Rightarrow g(3) = 7 \end{aligned}$$

$$\begin{aligned} g \circ f(2) &= g(f(2)) \\ &= g(3) \\ &= 7 \end{aligned}$$

e.g. Find  $g \circ f$  and  $f \circ g$ , if  $f: \mathbb{R} \rightarrow \mathbb{R}$  and  $g: \mathbb{R} \rightarrow \mathbb{R}$  are given by  $f(x) = \cos x$  and  $g(x) = 3x^2$ . Show that  $g \circ f \neq f \circ g$ .

$$\begin{aligned} g \circ f &= g \circ f(x) \\ &= g(f(x)) \\ &= g(\cos x) \\ &= 3(\cos x)^2 \\ &= 3 \cos^2 x \end{aligned}$$

$$\begin{aligned} f \circ g &= f \circ g(x) \\ &= f(g(x)) \\ &= f(3x^2) \\ &= \cos(3x^2) \end{aligned}$$

$g \circ f \neq f \circ g$

$x=0$

$3$

$x=0$

$1$




## Properties of Composite Functions

- ①  $g \circ f \neq f \circ g$  &  $h \circ (g \circ f) = (h \circ g) \circ f$
- ② If  $f$  and  $g$  are one-one, then  $g \circ f$  is one-one.
- ③ If  $f$  and  $g$  are onto, then  $g \circ f$  is onto.
- ④ If  $g \circ f$  is one-one, then 'f' is one-one.
- ⑤ If  $g \circ f$  is onto, then 'g' is onto.

## Invertible Functions (प्रतिलोमीय फलन) [ $f^{-1} / f^{-1}(x)$ ]

Eligibility / योग्यता :  $\rightarrow$  Bijective Function



one-one

&

onto

Let  $f(x_1) = f(x_2)$   
then  $x_1 = x_2$   
only

Range = Codomain

actually output

Question  
 $f: A \rightarrow B$

Definition: A function  $f: X \rightarrow Y$  is defined to be invertible, if there exists a function  $g: Y \rightarrow X$  such that  $g \circ f = I_X$  and  $f \circ g = I_Y$ .

The function 'g' is called the inverse of f.

$$g = f^{-1}$$

$$f \circ f^{-1} = I = f^{-1} \circ f$$

$$f \circ f^{-1}(x) = x = f^{-1} \circ f(x)$$

# Process to Find Inverse of a Function (f(x))

- check bijectivity of f(x).

f(x) = expression

$\swarrow$  one-one  $\searrow$   
onto

• Let  $y = f(x) = \boxed{\text{terms of } x}$

→ write  $x$  in terms of  $y$  ( $x = \boxed{\text{terms of } y}$ )

→ Replace  $x$  by  $f^{-1}(x)$


&  $y$  by  $x$ .

e.g. Find  $f^{-1}$  (if exists).

(i)  $S = \{1, 2, 3\}$

$f: S \rightarrow S$  given by

$f = \{(1, 3), (3, 2), (2, 1)\}$



$f(1) = 3$

$f(3) = 2$

$f(2) = 1 \Rightarrow f^{-1}(1) = 2$

one-one

onto ✓

Range = Codomain

$S = \{1, 2, 3\}$


(S)

$f^{-1} = \{(3, 1), (2, 3), (1, 2)\}$

(ii)  $S = \{1, 2, 3\}$

$f: S \rightarrow S$  given by

$f = \{(1, 2), (2, 1), (3, 1)\}$



one-one

X

→ not

bijjective

not invertible

e.g.  $f: \mathbb{R} \rightarrow \mathbb{R}$  defined by  $f(x) = 3x - 7$ . Find  $f^{-1}$  if exists.

Bijection

one-one

$f(x_1) = f(x_2)$

$\Rightarrow 3x_1 - 7 = 3x_2 - 7$

$\Rightarrow x_1 = x_2$  only

one-one

onto

Range = Codomain

$f: \mathbb{R} = \mathbb{R}$

$x \in \mathbb{R}$

$\Rightarrow 3x - 7 \in \mathbb{R}$

$\Rightarrow f(x) \in \mathbb{R}$

Range =  $\mathbb{R}$

onto

$$y = f(x) = 3x - 7 \quad (\text{let})$$

$$\Rightarrow y = 3x - 7$$

$$\Rightarrow \frac{y+7}{3} = x$$

$$\Rightarrow x = \frac{y+7}{3}$$

then (after replacement)

$$f^{-1}(y) = \frac{y+7}{3}$$


### Properties of Invertible Functions

①  $f \circ f^{-1}(x) = x = f^{-1} \circ f(x)$

②  $(f^{-1})^{-1} = f$

★ ③  $(g \circ f)^{-1} = f^{-1} \circ g^{-1}$

$$(g \circ f)^{-1}(x) = f^{-1} \circ g^{-1}(x)$$





Note: Let  $f: X \rightarrow Y$  is a bijective function.  
then  $f^{-1}: Y \rightarrow X$  is a bijective function.

Domain of  $f = X =$  Range of  $f^{-1}$

Range of  $f = Y =$  Domain of  $f^{-1}$

Graphically,  $f$  and  $f^{-1}$  are mirror image of each other in mirror line  $y=x$ .



mirror

$\log_e x$        $e^x$

inverse fun<sup>n</sup>.

Exercise 1.3 (Composite Fun<sup>n</sup> & Inverse Fun<sup>n</sup>)

Q.1  $f: \{1, 3, 4\} \rightarrow \{1, 2, 5\}$ ,  $g: \{1, 2, 5\} \rightarrow \{1, 3\}$  gof ≠ ?

$f = \{(1, 2), (3, 5), (4, 1)\}$ ,  $g = \{(1, 3), (2, 3), (5, 1)\}$

$f(1) = 2$ ,  $g(1) = 3$

$f(3) = 5$ ,  $g(2) = 3$



$f(4) = 1$ ,  $g(5) = 1$

$gof = \{(1, 3), (3, 1), (4, 3)\}$

$gof(1) = g(2) = 3$

$gof(3) = g(5) = 1$

$gof(4) = g(1) = 3$



$f \circ g = f \circ g(n) = f(g(n))$

Composite

Q.2  $f, g, h: R \rightarrow R$

Prove that: (i)  $(f+g) \circ h = f \circ h + g \circ h$

(ii)  $(f \cdot g) \circ h = (f \circ h) \cdot (g \circ h)$

$(f+g)(n) = f(n) + g(n)$

$(f \cdot g)(n) = f(n) \cdot g(n)$

$(f/g)(n) = \frac{f(n)}{g(n)}$

Algebra of  $F_n^m$

(ifm)

(i) LHS =  $(f+g) \circ h$   
 $= (f+g) \circ h(x)$

$= (f+g)(h(x))$

$= f(h(x)) + g(h(x))$

$= f \circ h + g \circ h$

= RHS

(ii) LHS =  $(f \cdot g) \circ h$

$= (f \cdot g)(h(x))$

$= f(h(x)) \cdot g(h(x))$

$= f \circ h \cdot g \circ h$

= RHS

③ Find gof and fog if

(i)  $f(x) = |x|$  and  $g(x) = |5x-2|$

$gof = g(f(x)) = g(|x|) = |5|x|-2|$  ✓

$|2| = |2| = 2$   
 $|-2| = |2| = 2$

$fog = f(g(x)) = f(|5x-2|) = ||5x-2| = |5x-2|$

(ii)  $f(x) = 8x^3$ ,  $g(x) = x^{1/3}$

$gof = gof(x) = g(f(x)) = g(8x^3) = (8x^3)^{1/3} = 8^{1/3} \cdot (x^3)^{1/3} = 2x$  ✓

$fog = f(g(x)) = f(x^{1/3}) = 8(x^{1/3})^3 = 8x$  ✓

④ If  $f(x) = \frac{4x+3}{6x-4}$ ,  $x \neq \frac{2}{3}$ , show that  $fof(x) = x$ ,  
 for all  $x \neq \frac{2}{3}$ . what is the inverse of 'f'?

To Prove:

$fof(x) = x$

LHS =  $fof(x)$

=  $f(f(x))$

=  $f\left(\frac{4x+3}{6x-4}\right)$

$fof(x) = \frac{4\left[\frac{4x+3}{6x-4}\right] + 3}{6\left[\frac{4x+3}{6x-4}\right] - 4}$

=  $\frac{6x + 12 + 18x - 12}{6x - 4}$   
 $= \frac{24x + 18 - 24x + 16}{6x - 4}$   
 $= \frac{34}{34} = x = \text{RHS}$

$f: x \rightarrow y$ ,  $f^{-1}: y \rightarrow x$   
 $f \circ f^{-1}(x) = x = f^{-1} \circ f(x)$   
 $f \circ g(x) = x = g \circ f(x)$

Let  
 $y = f(x) = \frac{4x+3}{6x-4}$

$$\Rightarrow y = \frac{4x+3}{6x-4}$$

$$\Rightarrow 6xy - 4y = 4x + 3$$

$$\Rightarrow 6xy - 4x = 4y + 3$$

$$\Rightarrow x(6y - 4) = 4y + 3$$

$$\Rightarrow x = \frac{4y+3}{6y-4}$$

$$\Rightarrow f^{-1}(x) = \frac{4x+3}{6x-4} = f(x)$$

① let  $y = f(x) = \dots$

②  $x$  in terms of  $y$

$$x = \dots$$


③  $x \rightarrow f^{-1}(x)$

$$y \rightarrow x$$

[Q.5] Invertible  $\begin{cases} \rightarrow \text{Yes} \\ \rightarrow \text{No} \end{cases}$  (with reason)

**Bijective**  
 $\swarrow \quad \searrow$   
 one-one onto


(i)  $f: \{1, 2, 3, 4\} \rightarrow \{10\}$ ,  $f = \{(1,10), (2,10), (3,10), (4,10)\}$



many-one

Not invertible


(ii)  $g: \{5, 6, 7, 8\} \rightarrow \{1, 2, 3, 4\}$ ,  $g = \{(5,4), (6,3), (7,4), (8,2)\}$



many-one & into

Not invertible

(iii)  $h: \{2, 3, 4, 5\} \rightarrow \{7, 9, 11, 13\}$ ,  $h = \{(2,7), (3,9), (4,11), (5,13)\}$



one-one, onto

Yes

Range = Codomain

Toppers Village (12<sup>th</sup> maths) (Relations & Functions)

Exercise 1.3 Composite Fun. & Inverse Fun.

Q.6 Show that  $f: [-1, 1] \rightarrow \mathbb{R}$ , given by  $f(x) = \frac{x}{x+2}$  is one-one.


Find the inverse of the function  $f: [-1, 1] \rightarrow$  Range of  $f$ .

Ans.  $f(x) = \frac{x}{x+2}$  one-one let  $f(x_1) = f(x_2) \Rightarrow$  only  $x_1 = x_2$

$$\Rightarrow \frac{x_1}{x_1+2} = \frac{x_2}{x_2+2}$$

$$\Rightarrow x_1 x_2 + 2x_1 = x_1 x_2 + 2x_2$$

$\Rightarrow$  only  $x_1 = x_2$   $\rightarrow f(x) \rightarrow$  one-one.



Inverse of  $f(x) = \frac{x}{x+2}$

let  $y = f(x) = \frac{x}{x+2}$

$$\Rightarrow y = \frac{x}{x+2}$$

$$\Rightarrow x = \frac{-2y}{y-1}$$

- ①  $y = f(x)$  (H1T1)
- ②  $x$  in terms of 'y'
- ③ Replace  $x \rightarrow f^{-1}(x)$   
 $y \rightarrow x$

$$\Rightarrow xy + 2y = x$$

$$\Rightarrow xy - x = -2y$$

$$\Rightarrow x(y-1) = -2y$$

$$\Rightarrow f^{-1}(x) = \frac{-2x}{x-1} = \frac{2x}{1-x}$$



Q.7 Consider  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by  $f(x) = 4x + 3$ . Show that  $f$  is invertible. Find the inverse of  $f$ .

Bijjective

one-one

let  $f(x_1) = f(x_2)$

$\Rightarrow 4x_1 + 3 = 4x_2 + 3$

$\Rightarrow x_1 = x_2$  only

one-one

✓

onto

Range = codomain

?

$\mathbb{R}$

(Actual output)

$x \rightarrow$  input  $x \in \mathbb{R}$   
 $f(x) \rightarrow$  output

$x \in \mathbb{R}$

$\Rightarrow 4x + 3 \in \mathbb{R}$

$\Rightarrow f(x) \in \mathbb{R}$

Range =  $\mathbb{R}$  = Codomain

onto ✓

$\therefore f$  is invertible,

$f(x) = 4x + 3 = y$  (let)

$\Rightarrow 4x = y - 3$

$\Rightarrow x = \frac{y - 3}{4}$

Replace  $x \rightarrow y \rightarrow f^{-1}(x)$

$y \rightarrow x$


$\Rightarrow f^{-1}(x) = \frac{x - 3}{4}$

$g \circ f(x) = x = f \circ g(x)$   
 $g \rightarrow f^{-1}$

Q.8 Consider  $f: \mathbb{R}_+ \rightarrow [4, \infty)$  given by  $f(x) = x^2 + 4$ .

Show that  $f$  is invertible with  $f^{-1}(y) = \sqrt{y-4}$ ,  
 where  $\mathbb{R}_+$  is the set of all non-negative real no.

(x) Domain =  $\mathbb{R}_+ = [0, \infty) \rightarrow x_1, x_2$   
 Codomain =  $[4, \infty)$



Invertible  $\rightarrow$  Bijective

one-one

onto

Let  $f(x_1) = f(x_2) \Rightarrow$  only  $x_1 = x_2$

Range = Codomain

Let  $f(x_1) = f(x_2)$

$$\Rightarrow x_1^2 + 4 = x_2^2 + 4$$

$$\Rightarrow x_1^2 = x_2^2$$

$$\Rightarrow x_1 = \pm x_2$$

$x_1 = x_2$

$x_1 = -x_2$

✓

x  
 $x_2 \rightarrow \mathbb{R}_+$

one-one

$x_1 \rightarrow \ominus$   
 X

f(x)  $\rightarrow x^2 + 4$

$$\because x \in [0, \infty)$$

$$\Rightarrow x^2 \in [0, \infty)$$

$$\Rightarrow x^2 + 4 \in [4, \infty)$$

$$f(x) \in [4, \infty)$$

$$\text{Range} = [4, \infty) = \text{Codomain}$$

onto ✓

Let  $y = f(x) = x^2 + 4$

$$\Rightarrow y = x^2 + 4$$

$$\Rightarrow y - 4 = x^2$$

$$\Rightarrow x = \pm \sqrt{y-4}$$

$$\because x \in \mathbb{R}_+ \Rightarrow x = \sqrt{y-4}$$

Replacement

$$f^{-1}(x) = \sqrt{x-4}$$

$$f^{-1}(100) = \sqrt{100-4}$$

$$f^{-1}(y) = \sqrt{y-4}$$

Q.9  $f: \mathbb{R}_+ \rightarrow [-5, \infty)$   $f(x) = 9x^2 + 6x - 5$

Show that  $f$  is invertible with  $f^{-1}(y) = \left( \frac{\sqrt{y+6} - 1}{3} \right)$ .

(x) Domain =  $\mathbb{R}_+ = [0, \infty)$

Codomain =  $[-5, \infty)$

Invertible  
↓  
Bijective.

One-one

Onto

Let  $f(x_1) = f(x_2) \Rightarrow \boxed{x_1 = x_2}$  (only)

$\Rightarrow 9x_1^2 + 6x_1 - 5 = 9x_2^2 + 6x_2 - 5$

$\Rightarrow 9(x_1^2 - x_2^2) + 6(x_1 - x_2) = 0$

$\Rightarrow 3(x_1 - x_2) \cdot [3(x_1 + x_2) + 2] = 0$

$x_1 = x_2$   
only  
Accept

$3(x_1 + x_2) + 2 = 0$

$\Rightarrow 3x_1 + 3x_2 + 2 = 0$

$\Rightarrow 3x_1 = -3x_2 - 2$

$\Rightarrow x_1 = -\frac{3x_2 + 2}{3}$

$x \in \mathbb{R}_+$   
 $x_1 \in \mathbb{R}_+$   
 $x_2 \in \mathbb{R}_+$   
Given.

$x_2 \rightarrow \oplus$   
 $x_1 \rightarrow \ominus$

Reject

One-one

Range = Codomain

$[-5, \infty)$

$f(x) = 9x^2 + 6x - 5$

→ Perfect Square.

$f(x) = 9x^2 + 6x - 5$

$= (3x)^2 + 2 \cdot (3x) \cdot 1 + 1^2 - 1 - 5$

$f(x) = (3x+1)^2 - 6$

$x \in [0, \infty)$

$\Rightarrow 3x \in [0, \infty)$

$\Rightarrow 3x+1 \in [1, \infty)$

$\Rightarrow (3x+1)^2 \in [1, \infty)$

$\Rightarrow \underline{(3x+1)^2 - 6} \in [-5, \infty)$

$f(x) \in [-5, \infty)$

Range =  $[-5, \infty) = \text{Codomain}$

onto

$$y = f(x) = 9x^2 + 6x - 5 = (3x+1)^2 - 6$$

①  $y = f(x)$

②  $x$  in terms  
of  $y$

$$y = (3x+1)^2 - 6$$

$$\Rightarrow y + 6 = (3x+1)^2$$

$$\Rightarrow \pm \sqrt{y+6} = 3x+1$$

$$\Rightarrow \pm \sqrt{y+6} - 1 = 3x$$

$$\Rightarrow x = \frac{\pm \sqrt{y+6} - 1}{3}$$

$\because x \in \mathbb{R}^+$

$$x = \frac{\sqrt{y+6} - 1}{3}$$

$$\Rightarrow f^{-1}(y) = \frac{\sqrt{y+6} - 1}{3}$$

$y = f(x)$   
 $f^{-1}(y) = x$

~~③  $x = f^{-1}(y)$~~

**Exercise 1.3** { Composite, Inverse Functions }


**Q.10** Let  $f: X \rightarrow Y$  be an invertible function.  
 Show that  $f$  has unique inverse.

Proof: (Proof by Contradiction)

Let  $f$  does not have a unique inverse

let  $\emptyset$  inverse of function  $f$  are  $g_1$  &  $g_2$ .

$g_1 \neq g_2$



$f \rightarrow$  invertible  
 Bijective.  
 one-one  $\swarrow$   
 onto  $\searrow$

$g_1: Y \rightarrow X$  also  $g_2: Y \rightarrow X$

$f(x_1) = f(x_2)$   
 $\Rightarrow x_1 = x_2$

$f \circ g_1(x) = I_Y = f \circ g_2(x) \leftarrow$  By Definition of Inverse.

$\Rightarrow f \circ g_1(x) = f \circ g_2(x)$

$\Rightarrow f(g_1(x)) = f(g_2(x)) \quad (\because f \rightarrow \text{one-one})$


$\Rightarrow g_1(x) = g_2(x)$

$\Rightarrow \underline{g_1 = g_2}$  This contradicts our assumption.

$\therefore f$  has a unique inverse.

Q.11  $f: \overset{A}{\{1,2,3\}} \rightarrow \overset{B}{\{a,b,c\}}$   $f(1)=a, f(2)=b, f(3)=c$ .

Find  $f^{-1}$ . Show that  $(f^{-1})^{-1} = f$



$$f = \{(1,a), (2,b), (3,c)\}$$

$$\therefore f^{-1} = \{(a,1), (b,2), (c,3)\} = g$$

$$(f^{-1})^{-1} = \{(1,a), (2,b), (3,c)\} = g^{-1}$$

$$\boxed{(f^{-1})^{-1} = f}$$

Q.12 Let  $f: X \rightarrow Y$  be an invertible function. Show that inverse of  $f^{-1}$  is  $f$  i.e.  $(f^{-1})^{-1} = f$ .

Ans.  $(f: X \rightarrow Y)$

Inverse of  $f = f^{-1} = g$  (let)  $\begin{pmatrix} f^{-1}: Y \rightarrow X \\ g: Y \rightarrow X \end{pmatrix}$

By definitions

$$g \circ f = \overset{f(x)=x}{I} = f \circ g$$

I  $\rightarrow$  Identity  
 $f \circ f^{-1} = f(x) = x$

$$g \circ f = I_X$$

$$f \circ g = I_Y$$

$$f \circ g = x$$



$$g \circ f = I_X$$

$$\Rightarrow g \circ f(x) = x$$

$$g \circ f = g \circ f(x)$$

$$\Rightarrow f(x) = g^{-1}(x)$$

$$\Rightarrow f(x) = (f^{-1})^{-1}(x)$$

$$f: X \rightarrow X$$

$$\Rightarrow f = (f^{-1})^{-1}$$

**Q.13** If  $f: \mathbb{R} \rightarrow \mathbb{R}$  be given by  $f(x) = (3-x^3)^{\frac{1}{3}}$ , then  $f \circ f(x)$  is

- (A)  $x^{\frac{1}{3}}$     (B)  $x^3$     (C)  $x$     (D)  $3-x^3$

Ans.  $f(x) = (3-x^3)^{\frac{1}{3}}$

$$f \circ f(x) = f(f(x)) = f\left(\underbrace{(3-x^3)^{\frac{1}{3}}}\right)$$

$$= \left(3 - \left[(3-x^3)^{\frac{1}{3}}\right]^3\right)^{\frac{1}{3}}$$

$$= \left[3 - (3-x^3)\right]^{\frac{1}{3}} = (3-3+x^3)^{\frac{1}{3}}$$

$$f \circ f(x) = x$$

**Q.14**  $f(x) = \frac{4x}{3x+4}$

$f^{-1} = g$

①  $y = f(x)$

②  $x \rightarrow y$  Terms

(A)  $g(y) = \frac{3y}{3-4y}$

~~(B)  $g(y) = \frac{4y}{4-3y}$~~

(C)  $g(y) = \frac{4y}{3-4y}$

(D)  $g(y) = \frac{3y}{4-3y}$

$$\Rightarrow y = f(x) = \frac{4x}{3x+4}$$

$$\Rightarrow y = \frac{4x}{3x+4}$$

$$\Rightarrow 3xy + 4y = 4x$$

$$\Rightarrow 4y = 4x - 3xy$$

$$\Rightarrow 4y = x(4-3y)$$

$$\Rightarrow x = \frac{4y}{4-3y}$$

$$\Rightarrow f^{-1}(y) = \frac{4y}{4-3y}$$

$$\Rightarrow g(y) = \frac{4y}{4-3y}$$

$y = f(x)$   
 $f^{-1}(y) = x$

Binary Operations [द्विआचारी संक्रियाएं]


For State Boards

Binary  $\rightarrow$  (2) दो

$100 + 35 + 98$   
~~100 + 35 + 98~~

$\oplus \ominus \otimes \oslash$

$3 \ 5 \ 7$





Set = Humans

Definition of Binary operations (\*) on A  $\rightarrow$  from A to A

A binary operation '\*' on a set A is a function  $* : A \times A \rightarrow A$ .

$* : A \times A \rightarrow A$

We denote  $*(a,b)$  by  $a*b$ .



eg.  $+(a,b) \ a,b \in \mathbb{N} \rightarrow +(a,b) = +(2,3) = 2+3 = 5$

$\underbrace{2,3}_{\mathbb{N}} \rightarrow \underbrace{2}_{\mathbb{N}} + \underbrace{3}_{\mathbb{N}} = 5$



e.g. Check, which of the following  $*$  is binary?

- (i) on  $\mathbb{Z}$ ,  $a * b = a - b$
- (ii) on  $\mathbb{N}$ ,  $a * b = a - b$
- (iii) on  $\mathbb{Q}$ ,  $a * b = a + (a+b)^2$
- (iv) on  $\mathbb{R} - \{-1\}$ ,  $a * b = \frac{a}{b+1}$

(i)  $\mathbb{Z} = \text{integers} = \{\dots, -2, -1, 0, 1, 2, \dots\}$   
 $a * b = a - b$       $a \in \mathbb{Z}, b \in \mathbb{Z}, a - b \in \mathbb{Z}$  ✓  
 $*$  → Binary ✓

(ii)  $\mathbb{N} = \{1, 2, 3, \dots\}$       $a \in \mathbb{N}, b \in \mathbb{N}, a - b \notin \mathbb{N}$   
 $a * b = a - b$       $a = 1, b = 5$   
 $1 * 5 = 1 - 5 = -4 \notin \mathbb{N}$      Not Binary

(iii)  $\mathbb{Q} = \text{Rational no.} = \text{परिमित सं.} = \frac{p}{q} \neq 0$   
 $a * b = a + (a+b)^2$ ,  $a \in \mathbb{Q}, b \in \mathbb{Q}, a + (a+b)^2 \in \mathbb{Q}$   
 Binary ✓

IV  $\mathbb{R} - \{-1\}$   
 2 inputs     1 output  
 $a * b = \frac{a}{b+1}$       $a \in \mathbb{R} - \{-1\}$  ✓  
 $b \in \mathbb{R} - \{-1\}$  ✓  
 $\frac{a}{b+1} \notin \mathbb{R} - \{-1\}$   
Not Binary

$a = -\frac{1}{2}, b = -\frac{1}{2}$   
 $(-\frac{1}{2}) * (-\frac{1}{2}) = \frac{-\frac{1}{2}}{-\frac{1}{2} + 1} = \frac{(-\frac{1}{2})}{(\frac{1}{2})} = -1$

# Commutativity (क्रम विनिमेयता)

$$\text{if } \underline{a * b = b * a}$$

Associativity (साहचर्यता) if  $\underline{(a * b) * c = a * (b * c)}$

Identity (तत्समक) =  $e$  (identity of  $*$ )  
 $a * e = \underline{a} = e * a$  (unique)

Inverse (प्रतिलोम) =  $b$  (मान)  
(Let)

$$\underline{a} * b = e = b * a$$

↓  
(inverse of  $a$ )

e.g. Consider a binary operation ' $*$ ' on the set  $\{1, 2, 3\}$  given by following operation table

(a)

$*$	1	2	3
1	<u>1</u>	2	3
2	2	<u>2</u>	3
3	3	3	<u>3</u>

$$\underline{a * b}$$

(iii) Is  $*$  commutative?

$$\underline{a * b = b * a}$$

$$\left\{ \begin{array}{l} 2 * 3 = 3 * 2 \\ 1 * 3 = 3 * 1 \\ 1 * 2 = 2 * 1 \end{array} \right.$$

(i) Compute  $(2 * 3) * 1$   
 $= (3) * 1 = 3$

(ii) Compute  $2 * (3 * 1)$   
 $= 2 * (3)$   
 $= 3$

e.g. Let  $*$  be a binary operation on the set  $\{1, 2, 3, 4, 5\}$  defined by

$$a * b = \text{HCF of } a \text{ \& } b.$$

- (i) Is  $*$  a binary operations?  
 (ii) Is  $*$  commutative?  
 (iii) find identity element for  $*$ ?

Ans.

$$\begin{array}{c}
 a * b = \text{HCF}(a, b) \\
 \swarrow \quad \downarrow \quad \downarrow \\
 A \quad A \quad A
 \end{array}$$

$$A = \{1, 2, 3, 4, 5\}$$

$$1 * 2 = \text{HCF}(1, 2) = 1 \in A$$

$$2 * 4 = \text{HCF}(2, 4) = 2 \in A$$

$$3 * 5 = \text{HCF}(3, 5) = 1 \in A$$

(i) Yes. (Binary  $\checkmark$ )

(ii)  $a * b = \text{HCF of } a \text{ \& } b = \text{HCF of } b \text{ \& } a$

Yes Commutative  $= b * a$

(iii) Identity element  $= e$

$$a * e = a = e * a$$

$$a \in A$$

$$a \in \{1, 2, 3, 4, 5\}$$

$$a = 5$$

$$5 * e = 5$$

$$\Rightarrow \text{HCF of } 5 \text{ \& } e = 5$$

$$\text{HCF } 5 \text{ \& } 5$$

$$e = 5$$

$$a = 4$$

$$4 * e = 4$$

$$\Rightarrow \text{HCF of } 4 \text{ \& } e = 4$$

$$e = 5 \quad e = 4$$

Not unique

$\Rightarrow$

$$e = 4$$

**Identity element does not exist**

## **All Useful Links:**

- **YouTube Channel : Toppers Village (Get full and free video lectures of Class 10 & 11 & 12 Maths) -**  
<https://www.youtube.com/toppersvillage>
- **Website :** [www.toppersvillage.com](http://www.toppersvillage.com)
- **Instagram :**  
<https://www.instagram.com/toppersvillage/>
- **Facebook Page:** [www.facebook.com/ToppersVillage/](http://www.facebook.com/ToppersVillage/)
- **Facebook Group:**  
<https://www.facebook.com/groups/ToppersVillage>