

# Toppers Village



## Continuity & Differentiability


Class Notes ( pdf )

**Class 12 Maths**

YouTube पर इस Chapter के FREE Videos देखिए ...  
(Playlist की Link नीचे है) ↓↓↓

[https://www.youtube.com/watch?v=RuG\\_63kYZLQ&list=PLPFrn0ppwwkS6SNnBuHQcxhSFu4z5XKZG](https://www.youtube.com/watch?v=RuG_63kYZLQ&list=PLPFrn0ppwwkS6SNnBuHQcxhSFu4z5XKZG)

In this Video

Limit  
(सीमा)Continuity  
(सततता)Limit of a function at  $x=a$  $f(x)$ Value of function in  
the neighbourhood of  
 $x=a$ . $(x=a \text{ के आस पास})$  $h = \text{very very small Positive real number}$  $\lim_{x \rightarrow a} f(x)$ Right Hand limit at  $x=a$  = RHL =  $\lim_{h \rightarrow 0} f(a+h) = f(a^+)$ Left hand limit at  $x=a$  = LHL =  $\lim_{h \rightarrow 0} f(a-h) = f(a^-)$ { limit of  
a function  
at  $x=a$  }# Limit exists at  $x=a$  if LHL = RHL = finite real no.



# How to Deal with Questions of 'Limit'?

Find  $\lim_{x \rightarrow a} f(x)$

$\lim_{x \rightarrow a} f(x)$  से ही काम चल जाता है

Normal Functions

$$\lim_{x \rightarrow 1} x^2 + 5$$

(Direct put)

$$= 1^2 + 5 = 6$$

$$\lim_{x \rightarrow 3} \frac{x^2 - 9}{x - 3}$$

$$= \frac{9 - 9}{3 - 3} = \frac{0}{0}$$

$$= \frac{(x-3)(x+3)}{(x-3)}$$

$$= \lim_{x \rightarrow 3} (x+3) = 6$$

$$f(x) = \begin{cases} \frac{\sin x}{x}, & x < 0 \\ x+1, & x \geq 0 \end{cases}$$

$$\lim_{x \rightarrow 0} f(x) \quad \frac{L}{0} \quad \frac{R}{0}$$

$$LHL = \lim_{h \rightarrow 0} f(0-h) = 1$$

$$RHL = 1$$

LHL, RHL अलग-अलग निकालना पड़ता है

Piecewise Functions  $[x], |x|, \operatorname{sgn}, \{x\}$

$$f(x) = \begin{cases} x-1, & x > 0 \\ 2x^2-1, & x \leq 0 \end{cases}$$

$$\lim_{x \rightarrow 0} f(x) = -1$$

$$LHL = -1$$

$$RHL = -1$$

Indeterminate Forms

$$\frac{0}{0}, \frac{\infty}{\infty}, \infty - \infty, 0 \cdot \infty, 1^\infty, 0^0, \infty^0$$

→ use Factorisation

→ use standard Forms

→ use L-Hospital Rule  
(only for  $\frac{0}{0}, \frac{\infty}{\infty}$ )

$$\lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1}$$

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$$

$$\lim_{x \rightarrow 0} \frac{\tan x}{x} = 1$$

etc.




# Continuity (सतता)

① Continuity at a Point ( $x=c$ )


- exclude (हटा)
- include (शामिल)

(सतत)  
Continuous


Discontinuous  
(असतत)



Discontinuous



Discontinuous



Continuous

# Continuous at Fixed Point → if we can draw the graph of the function around that point without lifting the pen from the plane of the paper.

# mathematically LHL =  $\lim_{x \rightarrow c^-} f(x)$  =  $f(c)$  = RHL

Continuous

$$\lim_{h \rightarrow 0} f(c-h) = f(c) = \lim_{h \rightarrow 0} f(c+h)$$

$$\lim_{x \rightarrow c} f(x) = f(c)$$

limit



Exact.



## ② Continuity in an Interval $[a, b]$

$(a, b)$

$f$  is said to be continuous in interval  $[a, b]$ , if it is continuous at every point in  $[a, b]$  including  $a$  &  $b$ .



Continuity at 'a' means

$$f(a) = \text{RHL}$$

$$f(a) = f(a^+)$$

Continuity at 'b' means

$$\text{LHL} = f(b)$$

$$f(b^-) = f(b)$$

# ★ A real function  $f$  is said to be continuous if it is continuous at every point in the domain of  $f$ .

"Draw graph without lifting pen in the domain"



$$f(x) = 2x + 3 \quad \text{✓}$$

$$f(x) = x^2 - x^3 + 1$$

$$f(x) = |x| \quad \text{Continuous}$$

$$f(x) = K \quad \text{✓}$$

$$f(x) = x \quad \text{✓}$$

$$f(x) = \sin x$$


$$f(x) = \frac{1}{x}, \quad x \neq 0$$

$$f(x) = \tan x, \quad x \neq (2n+1)\frac{\pi}{2}$$

$$f(x) = \frac{1}{\sin x}, \quad x \neq n\pi$$


$$f(x) = \begin{cases} x+2, & x < 0 \\ x, & x \geq 0 \end{cases}$$

Continuous




$$f(x) = \begin{cases} x+2, & x < 0 \\ x, & x \geq 0 \end{cases}$$

Discont.



$$f(x) = \begin{cases} x+2, & x < 0 \\ 1, & x = 0 \\ x, & x > 0 \end{cases}$$

Discont.




$$f(x) = [x] = \text{greatest integer function.}$$

$$[2] = 2 \quad [2.1] = 2$$

$$[2.9] = 2$$

$$[3] = 3$$

Discontinuous





# Important Points for Continuity

① Suppose  $f$  &  $g$  be two functions continuous at a real number  $x=c$ .

→  $f+g$  is continuous at  $x=c$   $(f(x)+g(x))$

→  $f-g$  is continuous at  $x=c$   $f(x)-g(x)$  ✓

→  $f \cdot g$  is continuous at  $x=c$   $f(x) \cdot g(x)$  ✓

→  $\frac{f}{g}$ ,  $g(c) \neq 0$  is continuous at  $x=c$ .  $\frac{f(x)}{g(x)}$  ✓

②  $f \circ g$  (or  $f(g(x))$ ) is continuous at  $x=c$  if

$g$  is continuous at  $x=c$  and

$f$  is continuous at  $x=g(c)$ .

(Composite Functions)

$f \circ g$

$f(x)$  input

$f(g(x))$

Continuity

$x=c$

$f(g(c))$

$g$  को  $x=c$  पर Continuous होना जरूरी है

$f$  को  $x=g(c)$  पर continuous होना जरूरी है

$$f(2) = 5$$

$$g(1) = 2$$

$$f(g(1)) = f(2) = 5$$



Exercise - 5.1 Chapter (5)

**Q.1** Prove that the function  $f(x) = 5x - 3$  is continuous at  $x=0$ , at  $x=-3$  and at  $x=5$ .

Ans.

$$f(x) = 5x - 3$$

$$\boxed{x=0}$$

$$f(0) = -3 \text{ (Exact)}$$

$$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} (5x - 3)$$

$$= 5(0) - 3$$

$$= -3$$

$$\therefore f(0) = \lim_{x \rightarrow 0} f(x)$$

$\therefore$  at  $x=0$ ,  $f$  is continuous.

$$\boxed{x=-3}$$

$$f(-3) = 5(-3) - 3$$

$$= -18$$

$$\lim_{x \rightarrow -3} f(x) = \lim_{x \rightarrow -3} (5x - 3)$$

$$= 5(-3) - 3$$

$$= -18$$

$$f(-3) = \lim_{x \rightarrow -3} f(x)$$

$\therefore$  at  $x=-3$ ,  $f$  is continuous.

$$\boxed{x=5}$$



$$\boxed{\text{Q.2}} \quad f(x) = 2x^2 - 1 \quad \text{at } x=3$$

$$f(3) = 2(3)^2 - 1 = 17$$

$$\lim_{x \rightarrow 3} f(x) = \lim_{x \rightarrow 3} (2x^2 - 1) = 17$$

$$\boxed{f(3) = \lim_{x \rightarrow 3} f(x)}$$

Continuous at  $x=3$



**Q.3** Examine the Continuity. Continuous Function

Continuous in their Domain

(a)  $f(x) = x - 5$

Domain =  $\mathbb{R} \leftarrow$  Real no.  
 $x \in \mathbb{R}$

Let  $\underline{x = c} \in \mathbb{R}$

$\underline{f(c) = c - 5}$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x - 5) = c - 5$$

$$\boxed{f(c) = \lim_{x \rightarrow c} f(x), \underline{\underline{c \in \mathbb{R}}}}$$

$\therefore f(x)$  is continuous fn.

(b)  $f(x) = \frac{1}{x-5}, x \neq 5$

Domain  $x - 5 \neq 0$

$\Rightarrow x \neq 5$

$\rightarrow x \in \mathbb{R} - \{5\}$

Let  $c \in \mathbb{R} - \{5\}$

$$\boxed{f(c) = \frac{1}{c-5}, c \in \mathbb{R} - \{5\}}$$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} \left( \frac{1}{x-5} \right) = \frac{1}{c-5}$$

$$\boxed{f(c) = \lim_{x \rightarrow c} f(x), c \in \mathbb{R} - \{5\}}$$

$\therefore f(x)$  is continuous Domain fn.



③  $f(x) = \frac{x^2 - 25}{x + 5}, x \neq -5$

Domain  $x + 5 \neq 0$

$\Rightarrow x \neq -5$

$x \in \mathbb{R} - \{-5\}$

Domain

$c \in \mathbb{R} - \{-5\}$

$f(c) = \frac{c^2 - 25}{c + 5}$  ✓

$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} \frac{x^2 - 25}{x + 5}$

$= \frac{c^2 - 25}{c + 5}$  ✓

$f(c) = \lim_{x \rightarrow c} f(x), c \in \mathbb{R} - \{-5\}$

Continuous f.m.

④  $f(x) = |x - 5|$

Domain  $= \mathbb{R}$

Let  $x = c \in \mathbb{R}$

$f(c) = |c - 5|$  ✓

$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} |x - 5|$

$= |c - 5|$  ✓

~~$f(x)$~~   $\lim_{x \rightarrow c}$

$f(c) = \lim_{x \rightarrow c} f(x), c \in \mathbb{R}$

Continuous f.m.

④  $f(x) = x^n$  at  $x = n$ ,  $n \in \text{Positive integer}$   $= \{1, 2, 3, \dots\}$   
 $n \in \mathbb{N}$

$f(n) = n^n$  ✓

$\lim_{x \rightarrow n} f(x) = \lim_{x \rightarrow n} x^n = n^n$  ✓


$f(n) = \lim_{x \rightarrow n} f(x), n \in \mathbb{N}$

Continuous f.m.



**Q.5** Is the function  $f$  defined by

$$f(x) = \begin{cases} x, & x \leq 1 \\ 5, & x > 1 \end{cases}; \text{ continuous at } x=0? \\ \text{at } x=1? \\ \text{at } x=2?$$



$x=1$  critical point

$x=1$

$$\lim_{x \rightarrow 1} f(x) \neq f(1)$$

$$\text{LHL} \neq \text{RHL} \neq f(1)$$

$$f(1^-) = f(1^+) = f(1)$$

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = f(1)$$

exact

$$\Rightarrow \lim_{x \rightarrow 1^-} [x] = \lim_{x \rightarrow 1^+} [5] = [1]$$

$$\Rightarrow 1 \neq 5 \neq 1$$

LHL

RHL

exact.

not continuous at  $x=1$

$x=0$  के लिए

$$f(x) = x$$

$$\lim_{x \rightarrow 0} f(x) = \underline{f(0)}$$

$$\Rightarrow 0 = 0$$

Continuous

$x=2$  के लिए

$$f(x) = 5$$

$$\lim_{x \rightarrow 2} f(x) = f(2)$$

$$\Rightarrow \lim_{x \rightarrow 2} 5 = 5$$

$$\Rightarrow 5 = 5$$

Continuous



Exercise-5.1Chapter-5

Find all points of discontinuity of  $f$ , where  $f$  is defined by —

[Q.6]  $f(x) = \begin{cases} 2x+3, & \text{if } x \leq 2 \\ 2x-3, & \text{if } x > 2 \end{cases}$

Piecewise Function( $x=2$  Critical Point) $x=2$  is Continuous

$$\text{LHL} \neq \text{RHL} \neq f(2)$$

$\uparrow$  Limit  $\quad \uparrow$  Exact  
 $\lim_{x \rightarrow 2} f(x)$

LHL  $\downarrow$   $2^-$   $\uparrow$  RHL  $2^+$

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x) = f(2) \quad \text{Exact}$$

$$\Rightarrow \lim_{x \rightarrow 2^-} (2x+3) = \lim_{x \rightarrow 2^+} (2x-3) = 2(2)+3$$

$$\Rightarrow 4+3 \neq 4-3 \neq 4+3$$

$$\Rightarrow 7 \neq 1 \neq 7$$

$\therefore$  Point of Discontinuity is  $x=2$ .

$$\begin{array}{c} 2x+3 \quad 2x-3 \\ \leftarrow \quad \rightarrow \\ \leftarrow \quad \rightarrow \end{array}$$

(2)

Dis.

Always Continuous.

Always Continuous.

 $x < 2$  $x > 2$



$$\boxed{\text{Q.7}} \quad f(x) = \begin{cases} |x|+3, & \text{if } x \leq -3 \\ -2x, & \text{if } -3 < x < 3 \\ 6x+2, & \text{if } x \geq 3 \end{cases}$$

we will check continuity at  $x = -3$  &  $x = 3$  only,  
and at rest of values of  $x$ , the function  
f(x) will be continuous.

$$\boxed{x = -3} \quad \text{LHL} = \text{RHL} = f(-3) \quad \text{at } x = -3$$

$$\lim_{x \rightarrow -3^-} (|x|+3) = \lim_{x \rightarrow -3^+} (-2x) = |-3|+3$$

$$\Rightarrow |-3|+3 = -2(-3) = 3+3$$

$$\Rightarrow \underline{6} = \underline{6} = \underline{6} \quad \text{at } x = -3$$

Continuous

$$\boxed{x = 3} \quad \text{LHL} \neq \text{RHL} = f(3) \quad \text{exact.} \quad \text{at } x = 3$$

$$\Rightarrow \lim_{x \rightarrow 3^-} (-2x) = \lim_{x \rightarrow 3^+} (6x+2) = (6(3)+2)$$

$$\Rightarrow -2(+3) = 6(3)+2 = 20$$

$$\Rightarrow \underline{-6} \neq 20 = 20$$

at  $x = 3$  discontinuous.

Point of Discontinuity =  $x = 3$



**Q.8**  $f(x) = \begin{cases} \frac{|x|}{x} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$


we have to check at  $x=0$ .

$LHL = RHL = f(0)$   
Exact.

$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = 0$

$\Rightarrow \lim_{x \rightarrow 0^-} \frac{|x|}{x} = \lim_{x \rightarrow 0^+} \frac{|x|}{x} = 0$

$\Rightarrow \frac{-x}{x} = \frac{+x}{x} = 0$



$\Rightarrow -1 \neq +1 \neq 0$   
 $\downarrow \quad \downarrow \quad \downarrow$   
 $LHL \neq RHL \neq f(0)$

$x=0 \notin \text{Discontinuous}$

**Q.9**  $f(x) = \begin{cases} \frac{x}{|x|} & \text{if } x < 0 \\ -1 & \text{if } x \geq 0 \end{cases}$

we have to check at  $x=0$ .

$LHL = RHL = f(0) \rightarrow$

$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = (-1)$   
Exact.

$\Rightarrow \lim_{x \rightarrow 0^-} \frac{x}{|x|} = \lim_{x \rightarrow 0^+} (-1) = (-1)$

$\Rightarrow \frac{x}{-x} = -1 = -1$

$\Rightarrow -1 = -1 = -1$

$LHL = RHL = f(0)$

Continuous at  $x=0$

$x \in \mathbb{R}$

~~no~~ Point of Discont.

**Q.10**  $f(x) = \begin{cases} x+1 & , x \geq 1 \\ x^2+1 & , x < 1 \end{cases}$

$x=1$

$x \in \mathbb{R} - \{1\} \rightarrow \text{cont.}$

check

$LHL = RHL = f(1)$

$\Rightarrow \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = (1+1)$

$\Rightarrow 1^2+1 = 1+1 = 2$

$\Rightarrow 2 = 2 = 2$

Cont.  $\rightarrow$  everywhere



**Q.11**  $f(x) = \begin{cases} x^3 - 3, & \text{if } x \leq 2 \\ x^2 + 1, & \text{if } x > 2 \end{cases}$

check at only  $x=2$

$$\text{LHL} = \lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} x^3 - 3$$

$$= 5$$

$$\text{RHL} = \lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} x^2 + 1$$

$$= 5$$

$$f(2) = 2^3 - 3 = 5$$

Continuous at  $x=2$

Continuous every where

**Q.12**

$$f(x) = \begin{cases} x^{10} - 1, & x \leq 1 \\ x^2, & x > 1 \end{cases}$$

~~check~~ check at only  $x=1$

$$\text{LHL} = \lim_{x \rightarrow 1^-} f(x)$$

$$= \lim_{x \rightarrow 1^-} (x^{10} - 1)$$

$$= 1 - 1 = 0$$

$$\text{RHL} = \lim_{x \rightarrow 1^+} (x^2)$$

$$= 1$$

$$f(1) = 1^{10} - 1 = 0$$

$$\text{LHL} = f(1) \neq \text{RHL}$$

$$\downarrow \quad \downarrow \quad \downarrow$$

Point of discontinuity

$x=1$

~~**Q.13**  $f(x) = \begin{cases} x+8, & \text{if } x \leq 5 \\ x-5, & \text{if } x > 5 \end{cases}$~~



Q.13

$$f(x) = \begin{cases} x+5, & x \leq 1 \\ x-5, & x > 1 \end{cases}$$

exact

$$f(1) = 1+5 = 6$$

Check at only  $x=1$

$$LHL = \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x+5) = 1+5 = 6$$

$$RHL = \lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (x-5) = 1-5 = -4$$

$$f(1) = LHL \neq RHL$$


Discontinuous at  $x=1$

Q.14

$$f(x) = \begin{cases} 3, & \text{if } 0 \leq x \leq 1 \\ 4, & \text{if } 1 < x < 3 \\ 5, & \text{if } 3 \leq x \leq 10 \end{cases}$$

for domain

check at  $x=0, 1, 3, 10$



$x=0$

$$f(0) = RHL \quad (x \rightarrow 0^+)$$

Closed  $\rightarrow [0, 10]$

$$\Rightarrow 3 = 3$$

Continuous

$x=1$

$$f(1) = LHL \neq RHL$$

$$\Rightarrow 3 = 3 \neq 4 \quad (\text{Discontinuous})$$

$x=3$

Discont.

$x=10$

$$\rightarrow LHL = f(10)$$

$$\Rightarrow 5 = 5$$

Continuous

Discontinuous



Q.15

$$f(x) = \begin{cases} 2x, & x < 0 \\ 0, & 0 \leq x \leq 1 \\ 4x, & x > 1 \end{cases}$$

Check at  $x=0, 1$

$x=0$  LHL = RHL =  $f(0)$   
 $(0^-)$   $(0^+)$

$$\Rightarrow 2(0) = (0) = 0$$

$$\Rightarrow 0 = 0 = 0$$

Continuous.

$x=1$  LHL  $\neq$  RHL  $\neq f(1)$   
 $x=(1^-)$   $x=(1^+)$   $x \in (1)$

$$\Rightarrow 0 \neq 4(1) \neq 0$$

$$0 \neq 4 \neq 0$$

Discontinuous.

Q.16

$$f(x) = \begin{cases} -2, & x \leq -1 \\ 2x, & -1 < x \leq 1 \\ 2, & x > 1 \end{cases}$$

Check at  $x=-1, 1$

$x=-1$  LHL = RHL =  $f(-1)$

$$\Rightarrow (-2) = 2(-1) = -2$$

$$\Rightarrow -2 = -2 = -2$$

Continuous.

$x=1$  LHL = RHL =  $f(1)$

$$2(1) = 2 = 2(1)$$

$$\Rightarrow 2 = 2 = 2$$

Continuous

Continuous.




Exercise 5.1 Chapter - 5

[Q.17] Find the relationship between ~~a~~ a and b so that the function f defined by

$$f(x) = \begin{cases} ax+1, & \text{if } x \leq 3 \\ bx+3, & \text{if } x > 3 \end{cases} \text{ is continuous at } x=3.$$

Ans. Given, Continuous at  $x=3$



$$\underline{LHL} = \underline{f(3)} = \underline{RHL}$$

$$\Rightarrow \lim_{x \rightarrow 3^-} (f(x)) = f(3) = \lim_{x \rightarrow 3^+} (f(x))$$

$$\lim_{x \rightarrow 3^-} (ax+1) = a(3)+1 = \lim_{x \rightarrow 3^+} (bx+3)$$

$$\Rightarrow \underline{3a+1} = \underline{3a+1} = \underline{3b+3} \Rightarrow 3a+1 = 3b+3$$

$$\Rightarrow \boxed{3a = 3b+2}$$

[Q.18] For what value of  $\lambda$ ,  $f(x) = \begin{cases} \lambda(x^2-2x), & x \leq 0 \\ 4x+1, & x > 0 \end{cases}$

is continuous at  $x=0$ ? what about continuity at  $x=1$ ?

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$$

$$\Rightarrow \lim_{x \rightarrow 0^-} \lambda(x^2-2x) = \lim_{x \rightarrow 0^+} (4x+1) = \lambda(x^2-2x)$$

$$\Rightarrow \boxed{0 \neq 1 \neq 0} \quad \text{no } \textcircled{\lambda}$$

$x=1$  के लिए.

$$f(x) = 4x+1$$

$$LHL = RHL = f(1)$$

Continuous.

$$\lambda \in \mathbb{R}$$



Q.19 Show that the function  $g(x) = x - [x]$  is discontinuous at all integral points.  
 ( $[x] \rightarrow$  greatest integer function)

Continuous

✓

LHL = f(c) = RHL

Domain =  $\mathbb{R}$  ( $x \in \mathbb{R}$ )

$g(x) = x - [x]$


Let integer  $(I) \in \mathbb{R}$  ( $x = I$ )

$$\begin{aligned} \text{LHL} &= \lim_{x \rightarrow I^-} g(x) \\ &= \lim_{x \rightarrow I^-} x - [x] \\ &= I^- - [I^-] \\ &= I - (I-1) \\ &= I - I + 1 = 1 \end{aligned}$$

$$\begin{aligned} \text{RHL} &= \lim_{x \rightarrow I^+} g(x) \\ &= \lim_{x \rightarrow I^+} (x - [x]) \\ &= I^+ - [I^+] \\ &= I - I = 0 \end{aligned}$$

$g(I) = x - [x] = I - [I] = I - I = 0$

$f(x) = [x]$  (Just before)



$$\begin{aligned} [1] &= 1 \\ [1.1] &= 1 \\ [1.999] &= 1 \\ [2] &= 2 \\ [2.1] &= 2 \\ [2.2] &= 2 \end{aligned}$$

$$\begin{aligned} [2.999] &= 2 \\ [3.0] &= 3 \\ [3.1] &= 3 \\ [3.2] &= 3 \end{aligned}$$

$$\begin{aligned} [3.9] &= 3 \\ [4] &= 4 \end{aligned}$$

LHL = 1, RHL = 0,  $g(I) = 0$

Discontinuous



Q.20. Is the function defined by  $f(x) = x^2 - \sin x + 5$  continuous at  $x = \pi$ ?

Ans.

$$LHL = RHL = f(\pi)$$

$$\lim_{x \rightarrow \pi} f(x) = f(\pi) \quad \text{(exact)}$$

$$\lim_{x \rightarrow \pi} f(x) = \lim_{x \rightarrow \pi} (x^2 - \sin x + 5)$$

$$= \pi^2 - \sin \pi + 5$$

$$= \pi^2 + 5$$

$$f(\pi) = \pi^2 - \sin \pi + 5$$

$$= \pi^2 - 0 + 5$$

$$= \pi^2 + 5$$

Yes

Q.21 Discuss the Continuity

We know that  $\sin x$ ,  $\cos x$  are continuous functions in their Domain ( $x \in \mathbb{R}$ ).

(a)  $f(x) = \sin x + \cos x$

(b)  $f(x) = \sin x - \cos x$

(c)  $f(x) = \sin x \cdot \cos x$

$\therefore \sin x + \cos x$ ,  $\sin x - \cos x$ ,  $\sin x \cdot \cos x$  are also continuous functions.

Q.22

Cosine, cosecant, secant, cotangent functions

$$\cos x$$

$$\csc x$$

$$\sec x$$

$$\cot x$$

Domain

$$\mathbb{R}$$

$$\mathbb{R} - \{n\pi\}$$

$$\mathbb{R} - \{(2m+1)\frac{\pi}{2}\}$$


$$\mathbb{R} - \{n\pi\}$$

$$x \in \mathbb{R} - \{n\pi\}$$

$$f(x) = \csc x$$

$$\lim_{x \rightarrow c} f(x) = \csc c$$

Continuity





Q.23 Find all points of discontinuity of  $f$ ,

where  $f(x) = \begin{cases} \frac{\sin x}{x}, & x < 0 \\ x+1, & x \geq 0 \end{cases}$

Ans. we have to check continuity at only  $x=0$ .

LHL =  $\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} \frac{\sin x}{x}$

Standard Form

Class 11  
 $\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$

$= 1$

RHL =  $\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (x+1) = 1$

$f(0) = (0)+1 = 1$   
 $x=0$

LHL = RHL = f(0)  
Continuous.

Continuous everywhere



Q.24

Determine if  $f$  defined by

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x} & , \text{ if } x \neq 0 \\ 0 & , \text{ if } x = 0 \end{cases} \text{ is a continuous function.}$$

$x=0$  at check.

$$f(0) = 0$$

$$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \left( x^2 \sin \frac{1}{x} \right)$$

$$= \sin \frac{1}{x} \in [-1, 1]$$

$$= (0)^2 \times (\text{Quantity in } [-1, 1])$$

$$= 0$$

$$\therefore \lim_{x \rightarrow 0} f(x) = f(0) = 0$$

$\therefore f(x)$  is a continuous function.

$$\left[ \begin{array}{c} \text{LHL} = \text{RHL} = f(0) ?? \\ \downarrow \quad \downarrow \\ \lim_{x \rightarrow 0^-} \quad \lim_{x \rightarrow 0^+} \\ \lim_{x \rightarrow 0} f(x) ? \end{array} \right]$$

$$\begin{aligned} & (0)^2 \cdot \sin\left(\frac{1}{0}\right) \\ & 0 \times \boxed{\sin(\infty)} \\ & 0 \times [-1, 1] = 0 \end{aligned}$$

$$\begin{aligned} & \frac{0}{0}, \frac{\infty}{\infty} \quad \boxed{0 \cdot \infty} \quad \boxed{\sin} \\ & \boxed{\frac{\sin x}{x}} \end{aligned}$$




Q.25 Examine the continuity of  $f$ , where

$$f \text{ is defined by } f(x) = \begin{cases} \sin x - \cos x, & x \neq 0 \\ -1, & x = 0 \end{cases}$$

~~lim~~ (Check continuity at  $x=0$  only.

{ at rest of the values of  $x$ ,  $f(x)$  will be continuous. }

$$LHL = RHL = f(0)$$



$$\lim_{x \rightarrow 0} f(x) = f(0)$$

→  $\boxed{-1 = -1}$

Continuous at  $x=0$

Continuous ~~at~~  
everywhere

$$f(0) = -1$$

$$\lim_{x \rightarrow 0} f(x)$$

$$= \lim_{x \rightarrow 0} (\sin x - \cos x)$$

$$= \sin 0 - \cos 0$$

$$= 0 - 1$$

$$= -1$$



Exercise (5.1) Chapter (5)

Find the value of 'K' so that 'f' is continuous at the indicated point.

Q.26

$$f(x) = \begin{cases} \frac{K \cos x}{\pi - 2x}, & \text{if } x \neq \frac{\pi}{2} \\ 3, & \text{if } x = \frac{\pi}{2} \end{cases}$$

at  $x = \frac{\pi}{2}$ ↑  
Continuous

$$\text{LHL} = \text{RHL} = f\left(\frac{\pi}{2}\right)$$

$$f\left(\frac{\pi}{2}\right) = 3$$

$$f\left(\frac{\pi}{2}\right) = f\left(\frac{\pi}{2}^+\right) = f\left(\frac{\pi}{2}^-\right)$$

$$\downarrow \quad \downarrow$$

$$\lim_{x \rightarrow \frac{\pi}{2}} f(x) = f\left(\frac{\pi}{2}\right)$$

$$\Rightarrow \lim_{x \rightarrow \frac{\pi}{2}} \frac{K \cos x}{\pi - 2x} = 3$$

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$$

$$\frac{K \cdot \cancel{\cos\left(\frac{\pi}{2}\right)} \rightarrow 0}{\pi - \cancel{2 \cdot \left(\frac{\pi}{2}\right)} \rightarrow 0} \left(\frac{0}{0} \text{ form}\right) \rightarrow \text{indeterminate Form}$$

$$\Rightarrow x \rightarrow \frac{\pi}{2} \quad x = \frac{\pi}{2} + h \quad \lim_{h \rightarrow 0} \quad \lim_{x \rightarrow \frac{\pi}{2}} \quad \lim_{\frac{\pi}{2} + h \rightarrow \frac{\pi}{2}} \quad \lim_{h \rightarrow 0}$$

Substitution

$$\Rightarrow \lim_{h \rightarrow 0} \frac{K \cdot \cos\left(\frac{\pi}{2} + h\right)}{\pi - 2\left(\frac{\pi}{2} + h\right)} = 3 \Rightarrow \lim_{h \rightarrow 0} \frac{-K \sin h}{\cancel{\pi} - \cancel{\pi} - 2h} = 3$$



$$\Rightarrow \lim_{h \rightarrow 0} \frac{k \sinh}{h} = 3$$

$$\Rightarrow \frac{k}{2} = 3$$

$$\Rightarrow \boxed{k=6}$$

Standard Form

$$\lim_{n \rightarrow 0} \frac{\sin n}{n} = 1$$

**[Q.27]**  $f(x) = \begin{cases} Kx^2, & x \leq 2 \\ 3, & x > 2 \end{cases}$  at  $x=2$ .

Continuous

$$\text{LHL} = f(2) = \text{RHL}$$

$$\Rightarrow f(2^-) = f(2) = f(2^+)$$

$$\lim_{x \rightarrow 2^-} Kx^2 = K(2)^2 = \lim_{x \rightarrow 2^+} (3)$$

$$\Rightarrow \underline{K(2)^2} = \underline{K(2)^2} = 3$$

$$\Rightarrow 4K = 3 \Rightarrow \boxed{K = \frac{3}{4}}$$

**[Q.28]**  $f(x) = \begin{cases} Kx+1, & x \leq \pi \\ \cos x, & x > \pi \end{cases}$  at  $x=\pi$ .

$$\text{LHL} = f(\pi) = \text{RHL}$$

$$\Rightarrow K(\pi)+1 = K(\pi)+1 = \cos(\pi)$$

$$\Rightarrow K\pi+1 = -1$$

$$\Rightarrow \boxed{K\pi = -2}$$

$$\boxed{K = \frac{-2}{\pi}}$$



**Q.29**  $f(x) = \begin{cases} Kx+1, & x \leq 5 \\ 3x-5, & x > 5 \end{cases}$  at  $x=5$  Continuous

$$LHL = f(5) = RHL$$

$$\Rightarrow (K(5)+1) = K(5)+1 = (3(5)-5)$$

$$\Rightarrow 5K+1 = 10 \Rightarrow K = \frac{9}{5}$$

**Q.30** Find the value of  $a$  &  $b$  such that the function  $f(x) = \begin{cases} 5, & x \leq 2 \\ ax+b, & 2 < x < 10 \\ 21, & x \geq 10 \end{cases}$  is a continuous function.

$x=2$

$$LHL = RHL = f(2)$$

$$\Rightarrow f(2^-) = f(2^+) = f(2)$$

$$\Rightarrow 5 = a(2) + b = 5$$

$$\Rightarrow 2a + b = 5 \quad \text{--- (1)}$$

$x=10$

$$LHL = RHL = f(10)$$

$$f(10^-) = f(10^+) = f(10)$$

$$\Rightarrow a(10) + b = 21 = 21$$

$$\Rightarrow 10a + b = 21 \quad \text{--- (2)}$$

$$eq^n (2) - eq^n (1)$$

$$10a + b = 21$$

$$2a + b = 5$$

$$-$$

$$8a = 16$$

$$a = 2$$

$$2a + b = 5$$

$$2(2) + b = 5$$

$$\Rightarrow b = 1$$




**Q.31** Show that  $f(x) = \cos(x^2)$  is a continuous function.

Concept - Composite Function  
(function in function)

$f(g(n))$   
 $g(n) \rightarrow \text{Continuous}$   $\mathbb{N} \in \mathbb{R}$   
 $f(n) \rightarrow \text{Continuous}$   $\mathbb{N} \in \mathbb{R}$   
 $\downarrow$   
 $f(g(n))$  or  $g(f(n))$   
 $\rightarrow \text{Continuous}$   $\mathbb{N} \in \mathbb{R}$

$f(x) = \cos(x^2)$       let  $g(x) = x^2 \rightarrow \text{Continuous}$   
 $\downarrow$   
 $h(x) = \underline{\cos x} \rightarrow \text{Continuous}$   
 $\Rightarrow h(g(x)) = \underline{\underline{\cos(x^2)}} \rightarrow \text{Continuous}$   
 $\therefore f(x) = \cos(x^2) \rightarrow \text{Continuous}$   
 $f_n^m$

Q.32 Show that  $f(x) = |\cos x|$  is a continuous f<sup>n</sup>.

Let  $g(x) = \cos x \rightarrow$  continuous   
 $h(x) = |x| \rightarrow$  continuous   
 $\Rightarrow h(g(x)) = |\cos x| \rightarrow$  continuous fn.  
fn.



Q. 33 Examine that  $\sin|x|$  is a continuous function.

$$\boxed{f(x) = \sin|x|}$$

$$\text{let } g(x) = |x| \rightarrow \text{continuous}$$

$$h(x) = \sin x \rightarrow \text{continuous}$$

$$h(g(x)) = \sin|x| \rightarrow \text{continuous}$$

Q. 34 Find all the points of discontinuity of  $f$  defined by  $f(x) = |x| - |x+1|$ .


Ans.

$$g(x) = |x| \rightarrow \text{continuous}$$



$x \in \mathbb{R}$

$$h(x) = |x+1| \rightarrow \text{continuous}$$



$x \in \mathbb{R}$

$$\underline{g(x) - h(x) = |x| - |x+1|}$$

Continuous  $\swarrow$

$f(x) = |x| - |x+1|$  is also continuous for  $x \in \mathbb{R}$ .

No points of discontinuity




Derivatives (अवकलन)


Rate of change  
(Daily life)

(Graph)

$$\text{Slope} = \frac{y_2 - y_1}{x_2 - x_1}$$



final - initial



$$\text{Derivative} = \text{Rate of change} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{(x+h) - (x)}$$

$$\text{Right Hand Derivative} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \text{RHD}$$

$$\text{Left Hand Derivative (LHD)} = \lim_{h \rightarrow 0} \frac{f(x) - f(x-h)}{(x) - (x-h)}$$

$$= \lim_{h \rightarrow 0} \frac{f(x) - f(x-h)}{x - x + h}$$

$$\text{LHD} = \lim_{h \rightarrow 0} \frac{f(x-h) - f(x)}{-h}$$



# A function  $f(x)$  is said to be differentiable at 'x' if (I) Continuous at x

(II)  $LHD = RHD$  (at x) = finite

$$\boxed{\lim_{h \rightarrow 0} \frac{f(x-h) - f(x)}{-h} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}}$$

(LHD) (RHD)

# Derivative of  $f(x) = y$

$$\hookrightarrow \frac{d(f(x))}{dx} = f'(x) = \frac{d(y)}{dx} = y' = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

↳ Diff. of  $f(x)$  w.r.t.  $x$  (y) First Rule

Standard Formulas

$$(1) \frac{d(x^n)}{dx} = n \cdot x^{n-1}$$

Constant  $= x^0$

$$\begin{aligned} &\rightarrow \frac{d(K)}{dx} = 0 \\ &\rightarrow \frac{d(x)}{dx} = 1 \\ &\rightarrow \frac{d(x^2)}{dx} = 2x \\ &\rightarrow \frac{d(\frac{1}{x})}{dx} = -\frac{1}{x^2} \\ &\rightarrow \frac{d(\sqrt{x})}{dx} = \frac{1}{2\sqrt{x}} \end{aligned}$$



## ② Trigonometric Functions.


$$\frac{d(\sin x)}{dx} = \cos x, \quad \frac{d(\cos x)}{dx} = -\sin x$$

$$\frac{d(\tan x)}{dx} = \sec^2 x, \quad \frac{d(\sec x)}{dx} = \sec x \cdot \tan x$$

$$\frac{d(\cot x)}{dx} = -\operatorname{cosec}^2 x, \quad \frac{d(\operatorname{cosec} x)}{dx} = -\operatorname{cosec} x \cdot \cot x$$

Rules ①  $(u+v)' = u' + v'$

$$\frac{d(u+v)}{dx} \nearrow$$



②  $(u-v)' = u' - v'$

③  $(u \cdot v)' = u' \cdot v + u \cdot v'$

Leibnitz Rule

$$(u \cdot v \cdot w)' = u' \cdot v \cdot w + u \cdot v' \cdot w + u \cdot v \cdot w'$$

④  $\left(\frac{u}{v}\right)' = \frac{u' \cdot v - u \cdot v'}{v^2}$

⑤  $(K \cdot u)' = K(u)'$

$$\frac{d(Ku)}{dx} = K \frac{d(u)}{dx}$$

(K = constant)



# Chain Rule (for composite function)

(function in function)

$$g(x) \text{ in } f(x) = f(g(x))$$

e.g.  $\sin(x^2) = g(p(x))$   
 $p(x) = x^2, g(x) = \sin x$

$$\sin(\tan(\cos x))$$

$$p(g(x)) = p(\sin x) \\ = (\sin x)^2 = \sin^2 x$$

Composite  
Fn.  
↓

$$\# \frac{d[f(g(x))]}{dx} = f'(g(x)) \times g'(x)$$

$$\# \frac{d[f(g(h(p(x))))]}{dx} = f'(g(h(p(x)))) \cdot g'(h(p(x))) \cdot h'(p(x)) \cdot p'(x)$$

e.g.  $f(x) = \sin(\tan(x^2))$

$$\frac{d(f(x))}{dx} = \frac{d(\sin(\tan(x^2)))}{dx}$$

fastest way.

$$= \cos(\tan(x^2)) \times \sec^2(x^2) \times 2x$$

II-way

$$\frac{d(\sin(\tan x^2))}{d(\tan x^2)} \cdot \frac{d(\tan x^2)}{d(x^2)} \cdot \frac{d(x^2)}{dx}$$



e.g.  $f(n) = \sqrt{\cot(2n+3)}$

$$\frac{d(f(n))}{dn} = \frac{d(\sqrt{\cot(2n+3)})}{dn}$$

$$= \left( \frac{1}{2\sqrt{\cot(2n+3)}} \right) \cdot (-\operatorname{cosec}^2(2n+3))$$

(2)

$$= \frac{-\operatorname{cosec}^2(2n+3)}{\sqrt{\cot(2n+3)}}$$

$$\frac{d(f(n))}{dn} = \frac{d\sqrt{\cot(2n+3)}}{\textcircled{dn}}$$

$$= \frac{d\sqrt{\cot(2n+3)}}{d(\cot(2n+3))} \cdot \frac{d(\cot(2n+3))}{d(2n+3)} \cdot \frac{d(2n+3)}{\textcircled{dn}}$$

$$= \frac{1}{2\sqrt{\cot(2n+3)}} \cdot (-\operatorname{cosec}^2(2n+3)) \cdot (2)$$

$$\sqrt{\cot(2n+3)}$$

$$\frac{d(\sqrt{x})}{dx} = \frac{1}{2\sqrt{x}}$$

$$\frac{d(\cot x)}{dx} = -\operatorname{cosec}^2 x$$

$$\frac{d(2n+3)}{dn}$$

$$= \frac{d(2n)}{dn} + \frac{d(3)}{dn}$$

$$= \frac{2(d(n))}{dn} + 0$$

$$= 2$$



## Exercise 5.2 chapter (5)

✓ Chain Rule  $\frac{d(f(g(x)))}{dx} = f'(g(x)) \cdot g'(x)$

✓  $(u \cdot v)' = u' \cdot v + u \cdot v'$

✓  $\left(\frac{u}{v}\right)' = \frac{u' \cdot v - u \cdot v'}{v^2}$

[Q.1] Differentiate the function with respect to 'x'.

$\sin(x^2+5)$

$\frac{d[\sin(x^2+5)]}{dx} = \cos(x^2+5) \cdot \frac{d(x^2+5)}{dx}$

(I)

$= \cos(x^2+5) \cdot (2x+0)$

$= 2x \cos(x^2+5)$

$\frac{d(x^n)}{dx} = n \cdot x^{n-1}$

(II)  $\frac{d(\sin(x^2+5))}{dx} = \frac{d(\sin(x^2+5))}{d(x^2+5)} \cdot \frac{d(x^2+5)}{dx}$

$= \cos(x^2+5) \cdot (2x+0)$



Q.2

 $\cos(\sin x)$ 

$$\frac{d(\cos x)}{dx} = -\sin x$$

$$\text{I} \quad \frac{d(\cos(\sin x))}{dx} = -\sin(\sin x) \cdot \cos x$$

$$\text{II} \quad \frac{d(\cos(\sin x))}{dx} = \frac{d(\cos(\sin x))}{d(\sin x)} \cdot \frac{d(\sin x)}{dx}$$

$$= -\sin(\sin x) \cdot \cos x$$

$$\text{Q.3} \quad \frac{d(\sin(ax+b))}{dx} = \cos(ax+b) \cdot (a \cdot 1 + 0)$$

$$= a \cdot \cos(ax+b)$$

$$\text{Q.4} \quad \frac{d(\sec(\tan \sqrt{x}))}{dx} = \sec(\tan \sqrt{x}) \cdot \tan(\tan \sqrt{x}) \cdot \sec^2(\sqrt{x}) \cdot \frac{1}{2\sqrt{x}}$$

$$\frac{d(\sec x)}{dx} = \sec x \cdot \tan x$$

$$\frac{d(x^{1/2})}{dx} = \frac{1}{2} \cdot x^{1/2-1}$$

$$\frac{d(\sec(\tan \sqrt{x}))}{dx} = \frac{d(\sec(\tan \sqrt{x}))}{d(\tan \sqrt{x})} \cdot \frac{d(\tan \sqrt{x})}{d(\sqrt{x})} \cdot \frac{d(\sqrt{x})}{dx}$$



Q.5

$$\frac{\sin(ant+b)}{\cos(cnt+d)}$$

$$\left(\frac{u}{v}\right)' = \frac{u'.v - u.v'}{v^2}$$

$$\frac{d}{dn} \left( \frac{\sin(ant+b)}{\cos(cnt+d)} \right)$$

$$= \frac{\frac{d}{dn}(\sin(ant+b)) \cdot \cos(cnt+d) - \sin(ant+b) \cdot \frac{d}{dn}(\cos(cnt+d))}{\cos^2(cnt+d)}$$

Chain Rule.      Chain

$$= \frac{\cos(ant+b) \cdot (a) \cdot \cos(cnt+d) - \sin(ant+b) \cdot (-\sin(cnt+d)) \cdot (c)}{\cos^2(cnt+d)}$$

$$= \frac{a \cdot \cos(ant+b) \cdot \cos(cnt+d) + c \cdot \sin(ant+b) \cdot \sin(cnt+d)}{\cos^2(cnt+d)}$$

Q.6

$$\cos x^3 \cdot \sin^2(x^5) = \underbrace{\cos(x^3)}_u \cdot \underbrace{[\sin(x^5)]^2}_v$$

$$(u \cdot v)' = u' \cdot v + u \cdot v'$$

$$\frac{d}{dn} [\cos x^3 \cdot (\sin x^5)^2] = \frac{d}{dn}(\cos x^3) \cdot (\sin x^5)^2 + (\cos x^3) \cdot \frac{d}{dn}[(\sin x^5)^2]$$

$$= \underbrace{-\sin(x^3) \cdot 3x^{3-1}}_{\text{Chain}} \cdot (\sin x^5)^2 + (\cos x^3) \cdot \underbrace{2(\sin x^5) \cdot \cos x^5}_{\text{Chain}} \cdot 5x^4$$

$$= -3x^2(\sin x^3) \cdot \sin^2(x^5) + 10x^4 \cdot \cos x^3 \cdot (\sin x^5) \cdot \cos x^5$$



Q.7

$$2 \sqrt{\cot(x^2)}$$

$$\left\{ \begin{aligned} \frac{d(\sqrt{x})}{dx} &= \frac{1}{2\sqrt{x}} \\ \frac{d(\cot x)}{dx} &= -\operatorname{cosec}^2(x) \\ \frac{d(x^2)}{dx} &= 2x \end{aligned} \right.$$

$$\begin{aligned} \frac{d(2 \sqrt{\cot x^2})}{dx} &= 2 \cdot \frac{1}{2 \sqrt{\cot x^2}} \cdot -\operatorname{cosec}^2(x^2) \cdot 2x \\ &= \frac{-2x \cdot \operatorname{cosec}^2 x^2}{\sqrt{\cot x^2}} \end{aligned}$$

Q.8

$$\cos \sqrt{x}$$

$$\frac{d(\cos \sqrt{x})}{dx} = -\sin \sqrt{x} \cdot \frac{1}{2\sqrt{x}} = \frac{-\sin \sqrt{x}}{2\sqrt{x}}$$


Q.9 Prove that the function  $f(x) = |x-1|$ ,  $x \in \mathbb{R}$  is not differentiable at  $x=1$ .

Solution

Condition for Differentiability at point  $x=1 \rightarrow$

(I) Continuous at  $x=1$  ✓

(II) At  $x=1$ ,  $LHD = RHD$



( $h > 0$ )


$$\lim_{h \rightarrow 0} \frac{f(1-h) - f(1)}{-h} = \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h}$$



$$f(x) = |x-1|$$

we know that

✓  $|x-1|$  is a continuous function in  $x \in \mathbb{R}$ .



$\therefore |x-1|$  is also continuous at  $x=1$ .

at  $x=1$  left Hand Derivative

$$\text{LHD} = \lim_{h \rightarrow 0} \frac{f(1-h) - f(1)}{-h}$$

$$= \lim_{h \rightarrow 0} \frac{|1-h-1| - |1-1|}{-h}$$

$$= \lim_{h \rightarrow 0} \frac{|-h| - 0}{-h}$$

$$= \lim_{h \rightarrow 0} \frac{|-h|}{-h}$$

$$= \lim_{h \rightarrow 0} \frac{h}{-h} = \textcircled{-1} = \text{LHD}$$

$$\begin{aligned} f(x) &= |x-1| \\ f(1-h) &= |(1-h)-1| \\ &= |-h| \\ &= -h \end{aligned}$$

$$\begin{aligned} |1-3| &= 3 = -(-3) \\ |-h| &= -(-h) = h \end{aligned}$$

$$\text{RHD} = \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{|1+h-1| - |1-1|}{h}$$

$$= \lim_{h \rightarrow 0} \frac{|h| - 0}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1 = \text{RHD}$$

$$f(x) = |x-1|$$

$\therefore \text{LHD} \neq \text{RHD} \Rightarrow \boxed{\text{not diff. at } x=1}$



**Q.10** Prove that the greatest integer function  $f(x) = [x]$ ,  $0 < x < 3$  is not differentiable at  $\underline{x=1}$  &  $\underline{x=2}$ .

$x=1$

(I) Continuity at  $x=1$

$$\lim_{x \rightarrow 1} f(x) = f(1)$$




$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^+} f(x) = f(1) \quad \text{(Exact)}$$

(LHL)

(RHL)

$$\Rightarrow \lim_{x \rightarrow 1^-} [x] = \lim_{x \rightarrow 1^+} [x] = [1]$$

$$\Rightarrow [1^-] = [1^+] = [1]$$



$$0 \neq 1$$


$$\text{LHL} \neq \text{RHL}$$

$\therefore f(x) = [x]$  is not continuous at  $x=1$

$\Rightarrow f(x) = [x]$  is not differentiable at  $x=1$ .

Similarly  $x=2$


$[x] \rightarrow$  piecewise fun<sup>n</sup>.



Prt.

at  $x=1$  &  $x=2$

$[x] =$  just floor



$$[2] = 2$$

$$[2.1] = 2$$

$$[2.2] = 2$$

$$[2.999] = 2$$

$$\downarrow$$
  

$$(3^-)$$



## Derivatives of

### Implicit Functions

### Inverse Trigonometric Fun<sup>n</sup>.

$$\begin{aligned} \rightarrow & x^2 - 3y^3 \sin x = 8 \\ \rightarrow & x - \sin(xy) + x^2 = y^2 \end{aligned}$$

$$(\sin^{-1}x, \cos^{-1}x, \dots)$$

Explicit Fun<sup>n</sup>.  $y = f(x)$

$$(\textcircled{y}) = \sin x, (\textcircled{y}) = x^2 - 3x + \cos x, (\textcircled{y}) = |x| - 3x^2$$

### Differentiation of Implicit Functions.

$$\left( \frac{dy}{dx} \right)$$

Start Diff. from one end to another

e.g.

$$x + \sin(xy) - y = 0$$

(Find derivative of  $y$  with respect to 'x')

find  $\frac{dy}{dx}$

{ Chain Rule  
u.v  
y/v }

$$x + \sin(xy) - y = 0$$

by Diff. w.r.t. 'x'

$$\frac{d(x)}{dx} = 1, \frac{d(y)}{dx}$$

$$\Rightarrow \frac{d(x)}{dx} + \frac{d(\sin(xy))}{dx} - \frac{d(y)}{dx} = 0$$

(Chain Rule)

$$\Rightarrow 1 + \cos(xy) \cdot \left\{ \frac{dx}{dx} \cdot y + x \cdot \frac{dy}{dx} \right\} - \frac{dy}{dx} = 0$$

$$\Rightarrow 1 + \cos(xy) \cdot \left\{ y + x \cdot \frac{dy}{dx} \right\} - \frac{dy}{dx} = 0$$



$$\Rightarrow 1 + \cos(xy) \cdot \left\{ y + x \left( \frac{dy}{dx} \right) \right\} - \left( \frac{dy}{dx} \right) = 0$$

$$\Rightarrow 1 + y \cdot \cos xy + x \cdot \cos(xy) \cdot \left( \frac{dy}{dx} \right) - \left( \frac{dy}{dx} \right) = 0$$

$$\Rightarrow \frac{dy}{dx} \cdot \{ x \cos(xy) - 1 \} = -(1 + y \cos xy)$$

$$\Rightarrow \frac{dy}{dx} = \frac{-(1 + y \cos xy)}{x \cos xy - 1} = \frac{(1 + y \cos xy)}{1 - x \cos xy}$$



# Derivatives of Inverse Trigonometric Functions

$$\frac{d(\sin^{-1}x)}{dx} = \frac{1}{\sqrt{1-x^2}}$$

$$\leftrightarrow \frac{d(\cos^{-1}x)}{dx} = \frac{-1}{\sqrt{1-x^2}}$$

$$\frac{d(\tan^{-1}x)}{dx} = \frac{1}{1+x^2}$$

$$\leftrightarrow \frac{d(\cot^{-1}x)}{dx} = \frac{-1}{1+x^2}$$

$$\frac{d(\sec^{-1}x)}{dx} = \frac{1}{|x|\sqrt{x^2-1}}$$

$$\frac{d(\csc^{-1}x)}{dx} = \frac{-1}{|x|\sqrt{x^2-1}}$$

e.g. Proof,  $y = \sin^{-1}x$   
(माना)

$$\Rightarrow \sin y = x$$

Diff. w. r. t. (x)

$$\frac{dy}{dx} = ?$$

$$\cos^2 y = 1 - \sin^2 y$$

$$\cos y = \sqrt{1 - \sin^2 y}$$

10th

$$\Rightarrow \frac{d(\sin y)}{dx} = \frac{dx}{dx} \Rightarrow \cos y \cdot \frac{dy}{dx} = 1$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{\cos y} = \frac{1}{\sqrt{1 - \sin^2 y}}$$

$$\Rightarrow \boxed{\frac{d(\sin^{-1}x)}{dx} = \frac{1}{\sqrt{1-x^2}}}$$

$$\sin^{-1}x + \cos^{-1}x = \frac{\pi}{2}$$

Property

$$\cos^{-1}x = \frac{\pi}{2} - \sin^{-1}x$$

Differentiate



★ e.g. Find  $\left(\frac{dy}{dx}\right)$  in  $y = \tan^{-1} \left( \frac{\sqrt{1+x^2} - 1}{x} \right)$   
V.V. Imp.

Solve  
 ↓  
 (आसानी) ~~Substitution~~  
 (Substitution) →  $\left(\frac{dy}{dx}\right) =$

$\sqrt{1-x^2} \rightarrow x = \sin \theta \rightarrow \sqrt{1-\sin^2 \theta} \rightarrow \cos \theta$   
 $\sqrt{1+x^2} \rightarrow x = \tan \theta \rightarrow \sqrt{1+\tan^2 \theta} \rightarrow \sec \theta$   
 $\sqrt{x^2-1} \rightarrow x = \sec \theta \rightarrow \sqrt{\sec^2 \theta - 1} \rightarrow \tan \theta$   
 (Standard Substitution)

$$y = \tan^{-1} \left( \frac{\sqrt{1+x^2} - 1}{x} \right)$$

replace  $x$  by  $\tan \theta$   $x = \tan \theta$

$$y = \tan^{-1} \left( \frac{\sqrt{1+\tan^2 \theta} - 1}{\tan \theta} \right) = \tan^{-1} \left( \frac{\sec \theta - 1}{\tan \theta} \right)$$

$$y = \tan^{-1} \left( \frac{\frac{1}{\cos \theta} - 1}{\frac{\sin \theta}{\cos \theta}} \right) = \tan^{-1} \left( \frac{\frac{1 - \cos \theta}{\cos \theta}}{\frac{\sin \theta}{\cos \theta}} \right)$$

$$y = \tan^{-1} \left( \frac{1 - \cos \theta}{\sin \theta} \right) = \tan^{-1} \left( \frac{\sin^2 \theta/2}{\sin \theta/2 \cos \theta/2} \right)$$



$$y = \tan^{-1} \left( \frac{\sin \theta/2}{\cos \theta/2} \right)$$

$$f^{-1}(f(x)) = x$$

$$y = \tan^{-1} \left( \tan \frac{\theta}{2} \right)$$

$$\Rightarrow y = \frac{\theta}{2}$$

$$x = \tan \theta \quad \text{H.T.}$$

$$\tan^{-1} x = \theta$$

$$y = \frac{\tan^{-1} x}{2}$$

$$\frac{dy}{dx} = \frac{d \left( \frac{\tan^{-1} x}{2} \right)}{dx} = \frac{1}{2} \frac{d(\tan^{-1} x)}{dx}$$

$$\boxed{\frac{dy}{dx} = \frac{1}{2} \cdot \frac{1}{1+x^2}}$$



Toppers Village Class 12 maths [Continuity & Differentiability]

Exercise - 5.3 Chapter (5)

[Q.1] Find  $\frac{dy}{dx}$ .

~~2x~~  $2x + 3y = \sin x$

Diff. w.r.t. 'x'

Direct Diff.  $\xrightarrow[\text{(u/v)}]{\text{(Chain) (u.v)}} \frac{dy}{dx}$  31/11/17

$$\Rightarrow \frac{d(2x)}{dx} + \frac{d(3y)}{dx} = \frac{d(\sin x)}{dx}$$

$$\Rightarrow 2 \times 1 + 3 \left( \frac{dy}{dx} \right) = \cos x$$

$$\Rightarrow \left[ \frac{dy}{dx} = \frac{\cos x - 2}{3} \right]$$

[Q.2]  $2x + 3y = \sin y$

by Diff. w.r.t. 'x'

$$\Rightarrow \frac{d(2x)}{dx} + \frac{d(3y)}{dx} = \frac{d(\sin y)}{dx}$$

$$\Rightarrow 2 + 3 \left( \frac{dy}{dx} \right) = \cos y \cdot \left( \frac{dy}{dx} \right)$$

$$\Rightarrow 2 = \frac{dy}{dx} (\cos y - 3)$$

$$\Rightarrow \left[ \frac{dy}{dx} = \frac{2}{\cos y - 3} \right]$$



**Q3**  $ax + by^2 = \cos y$

by diff. w.r.t. 'x'

$$\frac{d(x^2)}{dx} = 2x$$

$$\Rightarrow \frac{d(ax)}{dx} + \frac{d(by^2)}{dx} = \frac{d(\cos y)}{dx}$$

$$\Rightarrow ax + b \cdot 2y \cdot \frac{dy}{dx} = -\sin y \cdot \frac{dy}{dx}$$

$$\Rightarrow 2by \left( \frac{dy}{dx} \right) + \sin y \left( \frac{dy}{dx} \right) = -a$$

$$\Rightarrow \boxed{\frac{dy}{dx} = \frac{-a}{2by + \sin y}}$$

**Q.4**  $xy + y^2 = \tan x + y$

by diff. w.r.t. 'x'

$$\Rightarrow \frac{d(xy)}{dx} + \frac{d(y^2)}{dx} = \frac{d(\tan x)}{dx} + \frac{d(y)}{dx}$$

$$\boxed{(u \cdot v)' = u' \cdot v + u \cdot v'}$$

$$\Rightarrow \frac{d(x)}{dx} \cdot y + x \cdot \frac{dy}{dx} + 2y \cdot \frac{dy}{dx} = \sec^2 x + \frac{dy}{dx}$$

$$\Rightarrow x \left( \frac{dy}{dx} \right) + 2y \left( \frac{dy}{dx} \right) - \frac{dy}{dx} = \sec^2 x - y$$

$$\Rightarrow \frac{dy}{dx} = \frac{\sec^2 x - y}{x + 2y - 1}$$



**Q.5**  $x^2 + xy + y^2 = 100$

by diff. w.r.t. 'x'

$$\frac{d(C)}{dx} = 0$$

$$\Rightarrow \frac{d(x^2)}{dx} + \frac{d(xy)}{dx} + \frac{d(y^2)}{dx} = \frac{d(100)}{dx}$$

$$\Rightarrow \underline{2x} + \underline{1 \cdot y} + x \underline{\frac{dy}{dx}} + 2y \cdot \underline{\frac{dy}{dx}} = 0$$

$$\Rightarrow (x + 2y) \frac{dy}{dx} = -(2x + y)$$

$$\Rightarrow \boxed{\frac{dy}{dx} = -\frac{(2x+y)}{x+2y}}$$

$$\frac{d(x^n)}{dx} = n \cdot x^{n-1}$$

**Q.6**  $x^3 + x^2y + xy^2 + y^3 = 81$

by diff. w.r.t. (x)

$$\Rightarrow \frac{d(x^3)}{dx} + \frac{d(x^2y)}{dx} + \frac{d(xy^2)}{dx} + \frac{d(y^3)}{dx} = \frac{d(81)}{dx}$$

$$\Rightarrow 3x^2 + \left[ (2x) \cdot y + x^2 \cdot \underline{\frac{dy}{dx}} \right] + \left[ 1 \cdot y^2 + x \cdot 2y \cdot \underline{\frac{dy}{dx}} \right] + 3y^2 \cdot \underline{\frac{dy}{dx}} = 0$$

$\xrightarrow{(u \cdot v)'} \rightarrow u' \cdot v + u \cdot v'$

$$\Rightarrow \frac{dy}{dx} \cdot (x^2 + 2xy + 3y^2) = -(3x^2 + 2xy + y^2)$$

$$\boxed{\frac{dy}{dx} = -\frac{(3x^2 + 2xy + y^2)}{(x^2 + 2xy + 3y^2)}}$$



**Q.7**  $\sin^2 y + \cos xy = k \leftarrow \text{constant}$   
 by diff. w.r.t. 'x'

$$\begin{array}{c} (\sin y)^2 \\ \rightarrow \rightarrow \rightarrow \\ \textcircled{x^2} \\ \rightarrow \\ \textcircled{2x} \end{array}$$

$$\Rightarrow \frac{d(\sin^2 y)}{dx} + \frac{d(\cos(xy))}{dx} = \frac{d(k)}{dx}$$

chain                      chain, (u.v)

$$\Rightarrow 2 \sin y \cdot \cos y \cdot \frac{dy}{dx} + [-\sin(xy)] \cdot \frac{d(xy)}{dx} = 0$$

$$\Rightarrow 2 \sin y \cdot \cos y \cdot \frac{dy}{dx} - \sin(xy) \cdot \left\{ 1 \cdot y + x \cdot \frac{dy}{dx} \right\} = 0$$

11<sup>th</sup> class Formula

$$\Rightarrow \sin 2y \cdot \left( \frac{dy}{dx} \right) - y \sin xy - x \sin(xy) \cdot \left( \frac{dy}{dx} \right) = 0$$

$$\Rightarrow \frac{dy}{dx} (\sin 2y - x \sin xy) = y \sin xy$$

$$\Rightarrow \boxed{\frac{dy}{dx} = \frac{y \sin(xy)}{\sin 2y - x \sin xy}}$$



**[Q.8]**  $\sin^2 x + \cos^2 y = 1$   
by diff. w. r. t. 'x'

$\rightarrow (\sin x)^2$   
 $(\cos y)^2$

$(x^2) \rightarrow 2x$

$\Rightarrow \frac{d(\sin^2 x)}{dx} + \frac{d(\cos^2 y)}{dx} = \frac{d(1)}{dx}$

Chain Chain

$\Rightarrow 2 \sin x \cdot \cos x + 2 \cos y \cdot (-\sin y) \cdot \frac{dy}{dx} = 0$

Square Sin Square Cos y

$2 \sin \theta \cdot \cos \theta = \sin 2\theta$

$=$

$\neq$

$\Rightarrow \sin 2x = \sin 2y \cdot \frac{dy}{dx}$

$\Rightarrow \boxed{\frac{dy}{dx} = \frac{\sin 2x}{\sin 2y}}$



## Exercise-5.3

## Chapter 5

Q9 — Q15

ITF

Find  $\frac{dy}{dx}$ 

Q.9

$$y = \sin^{-1} \left( \frac{2x}{1+x^2} \right)$$

(Avoid  $\frac{\text{Chain Rule}}{x}$ )Trigonometric  
Substitution

↓ Solve

(आसान)

Diff.

$$y = \sin^{-1} \left( \frac{2x}{1+x^2} \right)$$

Put  $x = \tan \theta$  \*

$$\tan^{-1} x = \theta$$

$$y = \sin^{-1} \left( \frac{2 \tan \theta}{1 + \tan^2 \theta} \right) = \sin^{-1} (\sin 2\theta)$$

$$y = 2\theta$$

$$y = 2 \tan^{-1} x$$

Diff.

$$\frac{dy}{dx} = 2 \frac{d(\tan^{-1} x)}{dx} = \frac{2}{1+x^2}$$

Q.10

$$y = \tan^{-1} \left( \frac{3x-x^3}{1-3x^2} \right)$$

$$-\frac{1}{\sqrt{3}} < x < \frac{1}{\sqrt{3}}$$

Substitution

$$x = \tan \theta$$

$$\theta = \tan^{-1} x$$

$$y = \tan^{-1} \left( \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta} \right)$$

$$y = \tan^{-1} (\tan 3\theta)$$

$$y = 3\theta = 3 \tan^{-1} x$$

$$y = 3 \tan^{-1} x$$

Diff.

$$\frac{dy}{dx} = \frac{3}{1+x^2}$$



**Q.11**  $y = \cos^{-1} \left( \frac{1-x^2}{1+x^2} \right), \quad 0 < x < 1$

Substitution  $x = \tan \theta$

$\theta = \tan^{-1} x$

$$y = \cos^{-1} \left( \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \right)$$

$$y = \cos^{-1}(\cos 2\theta)$$

$$y = 2\theta$$

$$y = 2 \tan^{-1} x$$

$$\frac{dy}{dx} = 2 \times \frac{1}{1+x^2}$$

$$= \frac{2}{1+x^2}$$

**Q.12**  $y = \sin^{-1} \left( \frac{1-x^2}{1+x^2} \right), \quad 0 < x < 1$

Substitution  $x = \tan \theta$

$\theta = \tan^{-1} x$

$$y = \sin^{-1} \left( \frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \right)$$

$$y = \sin^{-1}(\cos 2\theta)$$

$$y = \sin^{-1} \left( \sin \left( \frac{\pi}{2} - 2\theta \right) \right)$$

$$y = \frac{\pi}{2} - 2\theta$$

$$y = \frac{\pi}{2} - 2 \tan^{-1} x$$

$$\frac{dy}{dx} = 0 - 2 \times \frac{1}{1+x^2} = \frac{-2}{1+x^2}$$

$\sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$



**Q.13**  $y = \cos^{-1} \left( \frac{2x}{1+x^2} \right), \quad -1 < x < 1$

Substitution,  $x = \tan \theta$

$\tan^{-1} x = \theta$

$$y = \cos^{-1} \left( \frac{2 \tan \theta}{1 + \tan^2 \theta} \right)$$

$$y = \cos^{-1} (\sin 2\theta)$$

$$y = \cos^{-1} \left( \cos \left( \frac{\pi}{2} - 2\theta \right) \right)$$

$$y = \frac{\pi}{2} - 2\theta$$

$$\boxed{y = \frac{\pi}{2} - 2 \tan^{-1} x}$$

Diff.

$$\frac{dy}{dx} = 0 - \frac{2}{1+x^2}$$

$$= -\frac{2}{1+x^2}$$

**Q.14**  $y = \sin^{-1} \left( 2x \sqrt{1-x^2} \right); \quad -\frac{1}{\sqrt{2}} < x < \frac{1}{\sqrt{2}}$

put  $x = \sin \theta$

$\sin^{-1} x = \theta$

$$y = \sin^{-1} \left( 2 \sin \theta \sqrt{1 - \sin^2 \theta} \right)$$

$$y = \sin^{-1} \left( \frac{2 \sin \theta \cos \theta}{1} \right)$$

$$y = \sin^{-1} (\sin 2\theta)$$

$$y = 2\theta$$

$$y = 2 \sin^{-1} x$$

Diff.

$$\frac{dy}{dx} = 2 \times \frac{1}{\sqrt{1-x^2}}$$



Q.15  $y = \sec^{-1}\left(\frac{1}{2x^2-1}\right), \quad 0 < x < \frac{1}{\sqrt{2}}$

put  $x = \cos \theta$   $\cos^{-1} x = \theta$

$$y = \sec^{-1}\left(\frac{1}{2\cos^2\theta-1}\right)$$

$$y = \sec^{-1}\left(\frac{1}{\cos 2\theta}\right) = \sec^{-1}(\sec 2\theta)$$

$$y = 2\theta = 2\cos^{-1}x$$

$$y = 2\cos^{-1}x$$

Diff  $\frac{dy}{dx} = 2 \times \left(\frac{-1}{\sqrt{1-x^2}}\right) = \frac{-2}{\sqrt{1-x^2}}$



Differentiation of logarithmic function & Exponential Fn.

$(a > 0)$   
 $(a \neq 1)$

$y = \log_a x$

$y = a^x$

$y = 2^x, 10^x$

(Common logarithm)

(Natural logarithm)

$(y = \log_{10} x)$

$y = \log_e(x) = \ln(x)$

P & C

M

~~log~~

$\log x$

$e = \text{Euler's no.}$   
 $e = 2.718 \dots$

$e$

$a$

$y = e^x \rightarrow \frac{dy}{dx} = e^x$

$\frac{d(a^x)}{dx} = a^x \cdot \log_e a$

$y = \log_e x \rightarrow \frac{dy}{dx} = \frac{1}{x}$

$\frac{d(\log_a x)}{dx} = \frac{1}{x \cdot \log_e a}$

$\frac{d(\log x)}{dx}$

e.g.  $y = 2^x \rightarrow \frac{dy}{dx} = 2^x \cdot \log_e 2$

$y = 10^x \rightarrow \frac{dy}{dx} = 10^x \cdot \log_e 10$

$y = e^x \rightarrow \frac{dy}{dx} = e^x$



e.g.  $y = \log x \rightarrow \left( \frac{dy}{dx} = \frac{1}{x} \right)$   
 (maths  $\log$  is Base = e)

e.g.  $y = \log_{10} x = \frac{1}{x \cdot \log_e 10}$

$\hookrightarrow y = \log_{10} x$

Base change theorem

$$\frac{\log_a x}{\log_a y} = \log_y x$$

$y = \frac{\log_e x}{\log_e 10}$  (Base change theorem)  
 $\log_e 10 \rightarrow \text{Constant.}$

Diff. —

$$\Rightarrow \frac{dy}{dx} = \frac{1}{\log_e 10} \cdot \frac{d(\log_e x)}{dx}$$

$$\left[ \frac{dy}{dx} = \frac{1}{\log_e 10} \cdot \frac{1}{x} \right]$$

e.g.  $y = \log(\log(x^2))$   
 (Chain Rule)

$$\frac{d(\log x)}{dx} = \frac{1}{x}$$

Diff.

$$\frac{dy}{dx} = \frac{1}{\log x^2} \cdot \frac{1}{x^2} \cdot (2x)$$

$$= \frac{2}{(\log x^2) \cdot x} \quad \checkmark$$



Toppers Village Class 12 Maths [Continuity and Differentiability]

Exercise 5.4 Chapter 5

$$\checkmark \quad \frac{d(\log x)}{dx} = \frac{d(\log_e x)}{dx} = \frac{d(\ln x)}{dx} = \frac{1}{x}$$

Differentiate  
with respect  
to 'x'  $\rightarrow$

$$\checkmark \quad \frac{d(e^x)}{dx} = e^x$$

$$\boxed{\text{Q.1}} \quad \frac{e^x}{\sin x} \left( \frac{u}{v} \right)' = \frac{u'v - u.v'}{v^2}$$

$$\frac{d \left( \frac{e^x}{\sin x} \right)}{dx} = \frac{e^x \cdot \sin x - e^x \cdot \cos x}{\sin^2 x} = \frac{e^x (\sin x - \cos x)}{\sin^2 x}$$

$$\boxed{\text{Q.2}} \quad e^{\sin^{-1} x} \quad (\text{fun. in fun.}) \rightarrow \text{Chain Rule}$$

$$\frac{d(e^{\sin^{-1} x})}{dx} = e^{\sin^{-1} x} \cdot \frac{1}{\sqrt{1-x^2}}$$

$$\boxed{\text{Q.3}} \quad \frac{d(e^{x^3})}{dx} = e^{x^3} \cdot 3x^2$$

$$\frac{d(x^3)}{dx} = 3 \cdot x^{3-1}$$

$$\boxed{\text{Q.4}} \quad \frac{d \left\{ \sin \left( \tan^{-1} (e^{-x}) \right) \right\}}{dx} = \cos \left( \tan^{-1} (e^{-x}) \right) \cdot \frac{1}{1 + (e^{-x})^2} \cdot e^{-x} \cdot (-1)$$

(Chain)



Q.5  $\log(\cos e^x)$  (Chain Rule)

$$\frac{d(\log(\cos e^x))}{dx} = \frac{1}{\cos(e^x)} \cdot (-\sin e^x) \cdot e^x$$

$$= -e^x \cdot \tan e^x$$

Q.6

$$e^x + e^{x^2} + \dots + e^{x^5}$$

$$\frac{d(x^n)}{dx} \rightarrow n \cdot x^{n-1}$$

$$\frac{d(e^x + e^{x^2} + e^{x^3} + e^{x^4} + e^{x^5})}{dx} = \frac{d(e^x)}{dx} + \frac{d(e^{x^2})}{dx} + \frac{d(e^{x^3})}{dx}$$

$$+ \frac{d(e^{x^4})}{dx} + \frac{d(e^{x^5})}{dx}$$

$$= e^x + e^{x^2} \cdot (2x) + e^{x^3} \cdot 3x^2 + e^{x^4} \cdot 4x^3$$

$$+ e^{x^5} \cdot 5x^4$$

Q.7

$$\sqrt{e^{\sqrt{x}}}$$

$$(e^{x^{1/2}})^{1/2}$$

$$(x^{1/2}) = \sqrt{x}$$

$$\frac{d(\sqrt{x})}{dx} = \frac{1}{2\sqrt{x}}$$

$$\frac{d(\sqrt{e^{\sqrt{x}}})}{dx} = \frac{1}{2\sqrt{e^{\sqrt{x}}}} \times e^{\sqrt{x}} \times \frac{1}{2\sqrt{x}}$$

$$= \frac{e^{\sqrt{x}}}{4\sqrt{x e^{\sqrt{x}}}}$$



Q.8

$\log(\log n)$

$(n > 1)$

$$\frac{d(\log x)}{dx} = \frac{1}{x}$$

$$\frac{d(\log(\log n))}{dn}$$

$$= \frac{1}{\log n} \cdot \frac{1}{n} = \frac{1}{n \log n}$$

Q.9

$$\frac{d\left(\frac{\cos x}{\log x}\right)}{dx}$$

$$= \frac{(-\sin x) \cdot \log x - \cos x \cdot \left(\frac{1}{x}\right)}{(\log x)^2}$$

$$\left(\frac{u}{v}\right)' = \frac{u' \cdot v - u \cdot v'}{v^2}$$

$$= - \left( \frac{x \sin x \cdot \log x + \cos x}{x (\log x)^2} \right)$$

Q.10

$$\frac{\cos(\log x + e^x)}{\text{Chain}}$$

$$\frac{d(\cos x)}{dx} = -\sin x$$

$$\frac{d(\cos(\log x + e^x))}{dx}$$

$$= -\sin(\log x + e^x) \cdot \left\{ \frac{1}{x} + e^x \right\}$$

$$= -\left(\frac{1}{x} + e^x\right) \cdot \sin(\log x + e^x)$$



Toppers Village class 12 maths [Continuity & Differentiability]


Logarithmic Differentiation (लघुगुणकीय अवकलन)

क्या? (i)  $\frac{\text{Variable}}{(\text{Variable})}$  (ii)  $\frac{(f_1(x))^{n_1} \cdot (f_2(x))^{n_2} \dots}{(g_1(x))^{k_1} \cdot (g_2(x))^{k_2} \dots}$

$x^n, (\sin x)^n, \tan x$

$\sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)}}$

कैसे?



$e = \text{Base}$

log की Properties: ①  $\log(m^n) = n \log m$

$\frac{d(\log x)}{dx} = \frac{1}{x}$

②  $\log(m \cdot n) = \log m + \log n$

③  $\log\left(\frac{m}{n}\right) = \log m - \log n$

e.g. Differentiate  $(\tan x)^x$  with respect to  $x$ .

(Var.)<sup>Var.</sup>

$\frac{dy}{dx} = ?$

$y = (\tan x)^x$

$\Rightarrow \log y = \log(\tan x)^x$

$\Rightarrow \log y = x \cdot \log(\tan x)$

$(u \cdot v)' = u' \cdot v + u \cdot v'$

by Diff. w.r.t. 'x'

$\Rightarrow \frac{1}{y} \cdot \left(\frac{dy}{dx}\right) = 1 \times \log \tan x + x \cdot \frac{\sec^2 x}{\tan x}$

$\Rightarrow \frac{dy}{dx} = (\tan x)^x \left( \log \tan x + x \cdot \frac{\sec^2 x}{\tan x} \right)$



e.g. Differentiate  $\sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)}}$  with respect to  $x$ .

$$\text{let } y = \sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)}} = \left( \frac{-}{-} \right)^{\frac{1}{2}}$$

(log)

$$\Rightarrow \log y = \log \left( \frac{(x-1)(x-2)}{(x-3)(x-4)} \right)^{\frac{1}{2}} = \frac{1}{2} \log \left( \frac{(x-1)(x-2)}{(x-3)(x-4)} \right)$$

$$\Rightarrow \log(y) = \frac{1}{2} \left\{ \log(x-1) + \log(x-2) - \log(x-3) - \log(x-4) \right\}$$

Diff. w.r.t.  $x \rightarrow$

$$\Rightarrow \frac{1}{y} \cdot \left( \frac{dy}{dx} \right) = \frac{1}{2} \left\{ \frac{1}{x-1} \cdot x(1-0) + \frac{1}{(x-2)} - \frac{1}{x-3} - \frac{1}{(x-4)} \right\}$$

$$\Rightarrow \frac{dy}{dx} = \sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)}} \cdot \frac{1}{2} \cdot \left\{ \frac{1}{x-1} + \frac{1}{x-2} - \frac{1}{x-3} - \frac{1}{x-4} \right\}$$



Exercise-5.5 Chapter-5Logarithmic Differentiation① Variable  
(Variable)②  
 $(f_1(x))^{n_1} (f_2(x))^{n_2} \dots$   
 $(g_1(x))^{k_1} (g_2(x))^{k_2} \dots$ 

①  $y = \square$

②  $\log$

③  $\log$  Properties

$\log(m^n) = n \log m$

$\log(m \cdot n) = \log m + \log n$

$\log\left(\frac{m}{n}\right) = \log m - \log n$

④ Diff.

$\left(\frac{dy}{dx}\right)$

Exercise (5.5)

$(u \cdot v)' = u'v + u \cdot v'$

[Q.1] Let  $y = \cos x \cdot \cos 2x \cdot \cos 3x$

$\Rightarrow \log y = \log(\cos x \cdot \cos 2x \cdot \cos 3x)$

$\Rightarrow \log y = \log \cos x + \log \cos 2x + \log \cos 3x$

by Differentiating w.r.t. 'x'

$\frac{d(\log x)}{dx} = \frac{1}{x}$

$$\Rightarrow \frac{1}{y} \cdot \left(\frac{dy}{dx}\right) = \frac{1}{\cos x} \cdot (-\sin x) + \frac{1}{\cos 2x} \cdot (-\sin 2x) \cdot (2) + \frac{1}{\cos 3x} \cdot (-\sin 3x) \cdot (3)$$

$$\Rightarrow \frac{dy}{dx} = y \cdot (-\tan x - 2 \tan 2x - 3 \tan 3x)$$

$$= -\cos x \cos 2x \cos 3x (\tan x + 2 \tan 2x + 3 \tan 3x)$$



Q.2

$$y = \sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)}} = \left( \frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)} \right)^{1/2}$$

$$\Rightarrow \log y = \log \left[ \frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)} \right]^{1/2}$$

$$\Rightarrow \log y = \frac{1}{2} \cdot \log \left\{ \frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)} \right\}$$

$$\Rightarrow \log y = \frac{1}{2} \left\{ \log(x-1) + \log(x-2) - \log(x-3) - \log(x-4) - \log(x-5) \right\}$$

Diff. w.r.t. 'x'

$$\Rightarrow \frac{1}{y} \cdot \left( \frac{dy}{dx} \right) = \frac{1}{2} \left\{ \frac{1}{x-1} \cdot (1-0) + \frac{1}{x-2} - \frac{1}{x-3} - \frac{1}{x-4} - \frac{1}{x-5} \right\}$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{2} \sqrt{\frac{(x-1)(x-2)}{(x-3)(x-4)(x-5)}} \cdot \left\{ \frac{1}{x-1} + \frac{1}{x-2} - \frac{1}{x-3} - \frac{1}{x-4} - \frac{1}{x-5} \right\}$$

Q.3  $y = (\log x)^{\cos x}$

$$\Rightarrow \log y = \log [(\log x)^{\cos x}]$$

$$\Rightarrow \log y = \frac{\cos x \cdot \log(\log x)}{(u \cdot v) = u'v + u \cdot v'}$$

Diff. w.r.t. 'x'

$$\Rightarrow \frac{1}{y} \cdot \left( \frac{dy}{dx} \right) = -\sin x \cdot (\log \log x) + \cos x \cdot \frac{1}{\log x} \cdot \frac{1}{x}$$

$y = (\log x)^{\cos x}$

$$\frac{dy}{dx} = (\log x)^{\cos x} \cdot \left\{ -\sin x \cdot \log \log x + \frac{\cos x}{x \log x} \right\}$$



**Q.4**

$$y = \underset{\substack{\downarrow \\ u}}{x^n} - \underset{\substack{\downarrow \\ v}}{2^{\sin x}}$$

$$\Rightarrow y = u - v$$

$$\hookrightarrow \frac{dy}{dx} = \frac{du}{dx} - \frac{dv}{dx}$$

$$u = x^n$$

$$\Rightarrow \log u = \log(x^n)$$

$$\Rightarrow \log u = \underline{n \cdot \log x}$$

Diff. w.r.t. 'x'

$$\Rightarrow \frac{1}{u} \cdot \frac{du}{dx} = 1 \cdot \log x + x \cdot \frac{1}{x}$$

$$\Rightarrow \frac{du}{dx} = u(\log x + 1)$$

$$\Rightarrow \frac{du}{dx} = x^n (\log x + 1)$$

$$v = 2^{\sin x}$$

$$\Rightarrow \log v = \log(2^{\sin x})$$

$$\Rightarrow \log v = \sin x \cdot \log 2$$

Diff. w.r.t. 'x' Constant

$$\Rightarrow \frac{1}{v} \cdot \frac{dv}{dx} = \log 2 \cdot \cos x$$

$$\Rightarrow \frac{dv}{dx} = 2^{\sin x} \cdot \log 2 \cdot \cos x$$

$$\therefore \frac{dy}{dx} = \frac{du}{dx} - \frac{dv}{dx}$$

$$\Rightarrow \frac{dy}{dx} = \underline{x^n (\log x + 1)} - \underline{2^{\sin x} \cdot \log 2 \cdot \cos x}$$

**Q.5**

$$y = (x+3)^2 \cdot (x+4)^3 \cdot (x+5)^4$$

$$\Rightarrow \log y = \log((x+3)^2 \cdot (x+4)^3 \cdot (x+5)^4)$$

$$\Rightarrow \log y = 2 \log(x+3) + 3 \log(x+4) + 4 \log(x+5)$$

Diff. w.r.t. 'x'

$$\Rightarrow \frac{1}{y} \cdot \left( \frac{dy}{dx} \right) = \frac{2}{x+3} + \frac{3}{x+4} + \frac{4}{x+5}$$

$$\Rightarrow \frac{dy}{dx} = (x+3)^2 (x+4)^3 (x+5)^4 \left( \frac{2}{x+3} + \frac{3}{x+4} + \frac{4}{x+5} \right)$$



$$\Rightarrow \frac{dy}{dx} = (x+3)^1 (x+4)^2 (x+5)^3 \cdot \left\{ \begin{aligned} & \frac{2}{\cancel{x+3}} (\cancel{x+3}) (\cancel{x+4}) (\cancel{x+5}) \\ & + \frac{3}{\cancel{x+4}} (\cancel{x+3}) (\cancel{x+4}) (\cancel{x+5}) \\ & + \frac{4}{\cancel{x+5}} (\cancel{x+3}) (\cancel{x+4}) (\cancel{x+5}) \end{aligned} \right\}$$

$$\Rightarrow \frac{dy}{dx} = (x+3)^1 (x+4)^2 (x+5)^3 \cdot \left\{ \begin{aligned} & 2(x^2 + 9x + 20) \\ & + 3(x^2 + 8x + 15) \\ & + 4(x^2 + 7x + 12) \end{aligned} \right\}$$

$$\frac{dy}{dx} = (x+3)^1 (x+4)^2 (x+5)^3 \cdot \{ 9x^2 + 70x + 133 \}$$


---



Exercise 5.5 Chapter-5

Q.6  $\left(x + \frac{1}{x}\right)^x + x^{(1 + \frac{1}{x})}$

$y = \left(x + \frac{1}{x}\right)^x + x^{(1 + \frac{1}{x})}$

$y = u + v$

$\frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$

$u = \left(x + \frac{1}{x}\right)^x$

$\Rightarrow \log u = \log \left(x + \frac{1}{x}\right)^x$

$\Rightarrow \log u = x \cdot \log \left(x + \frac{1}{x}\right)$

Diff. w.r.t.  $x$

$\Rightarrow \frac{1}{u} \cdot \left(\frac{du}{dx}\right) = 1 \times \log \left(x + \frac{1}{x}\right) + x \cdot \frac{1}{x + \frac{1}{x}} \cdot \left(1 - \frac{1}{x^2}\right)$

$\Rightarrow \frac{du}{dx} = u \left[ \log \left(x + \frac{1}{x}\right) + \frac{x \cdot x}{x^2 + 1} \cdot \left(\frac{x^2 - 1}{x^2}\right) \right]$

$\Rightarrow \frac{du}{dx} = \left(x + \frac{1}{x}\right)^x \cdot \left[ \log \left(x + \frac{1}{x}\right) + \frac{x^2 - 1}{x^2 + 1} \right]$

$\log(m^n)$   
 $\downarrow$   
 $n \cdot \log m$   
Diff. -  $\frac{dy}{dx}$

$v = x^{(1 + \frac{1}{x})}$

$\log v = \left(1 + \frac{1}{x}\right) \cdot \log x$

Diff. w.r.t.  $x$

$\Rightarrow \frac{1}{v} \cdot \frac{dv}{dx} = \left(0 - \frac{1}{x^2}\right) \cdot \log x + \left(1 + \frac{1}{x}\right) \cdot \frac{1}{x}$

$\Rightarrow \frac{dv}{dx} = v \left[ \frac{-\log x}{x^2} + \frac{n+1}{x^2} \right]$

$\Rightarrow \frac{dv}{dx} = \frac{x^{(1 + \frac{1}{x})}}{x^2} \left[ -\log x + \frac{n+1}{x} \right]$



$$\therefore \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$$

$$\Rightarrow \frac{dy}{dx} = \left(x + \frac{1}{x}\right)^x \left[ \log\left(x + \frac{1}{x}\right) + \frac{x^2 - 1}{x^2 + 1} \right] + \frac{x^{(1 + \frac{1}{x})}}{x^2} \left[ -\log x + x + 1 \right]$$

**Q. 7**  $y = (\log x)^x + x^{\log x}$

$y = u + v$

$$\left[ \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx} \right]$$

$u = (\log x)^x$

$\Rightarrow \log u = x \cdot \log(\log x)$

Diff. w.r.t. (x)

$\Rightarrow \frac{1}{u} \cdot \left(\frac{du}{dx}\right) = 1 \cdot \log(\log x) + x \cdot \frac{1}{\log x} \cdot \frac{1}{x}$

$\Rightarrow \frac{du}{dx} = (\log x)^x \cdot \left[ \log \log x + \frac{1}{\log x} \right]$

$v = x^{\log x}$

$\log v = \log \left( x^{\log x} \right)$

$\Rightarrow \log v = \log x \cdot \log x$

$\Rightarrow \log v = (\log x)^2$

Diff. w.r.t. (x)

$\Rightarrow \frac{1}{v} \cdot \left(\frac{dv}{dx}\right) = 2 \log x \cdot \frac{1}{x}$

$\Rightarrow \frac{dv}{dx} = x^{\log x} \cdot \frac{2 \log x}{x}$

$\therefore \frac{dy}{dx} = (\log x)^x \cdot \left( \log \log x + \frac{1}{\log x} \right) + x^{\log x} \cdot \frac{2 \log x}{x}$



$$\boxed{\text{Q.8}} \quad y = (\sin x)^x + \sin^{-1}(\sqrt{x})$$

$$y = u + v$$

$$\boxed{\frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}}$$

$$u = (\sin x)^x$$

$$\Rightarrow \log u = \underline{x \cdot \log(\sin x)}$$

Diff. w.r.t. x

$$\Rightarrow \frac{1}{u} \cdot \frac{du}{dx} = 1 \cdot \log(\sin x) + x \cdot \frac{1}{\sin x} \cdot \cos x$$

$$\Rightarrow \frac{du}{dx} = (\sin x)^x \cdot [\log \sin x + x \cdot \cot x]$$

$$\therefore \frac{dy}{dx} = (\sin x)^x \cdot [\log \sin x + x \cdot \cot x] + \frac{1}{2\sqrt{x-x^2}}$$

$$v = \sin^{-1}(\sqrt{x})$$

chain rule

$$\Rightarrow \frac{dv}{dx} = \frac{1}{\sqrt{1-x}} \cdot \frac{1}{2\sqrt{x}}$$

$$\Rightarrow \frac{dv}{dx} = \frac{1}{2\sqrt{(1-x) \cdot x}}$$

$$\frac{dv}{dx} = \frac{1}{2\sqrt{x-x^2}}$$

$$\boxed{\text{Q.9}} \quad y = \underbrace{x^{\sin x}}_u + \underbrace{(\sin x)^{\cos x}}_v$$

$$\rightarrow y = u + v$$

$$\hookrightarrow \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$$



$$u = x^{\sin x}$$

$$\Rightarrow \log u = \log (x^{\sin x})$$

$$\Rightarrow \log u = \sin x \cdot \log x$$

Diff. w. r. t.  $(x)$

$$\Rightarrow \frac{1}{u} \cdot \frac{du}{dx} = \cos x \cdot \log x + \sin x \cdot \frac{1}{x}$$

$$\Rightarrow \frac{du}{dx} = (x^{\sin x}) \cdot \left[ \cos x \cdot \log x + \frac{\sin x}{x} \right] \checkmark$$

$$v = (\sin x)^{\cos x}$$

$$\log v = \cos x \cdot \log (\sin x)$$

Diff.  $(x)$

$$\Rightarrow \frac{1}{v} \cdot \frac{dv}{dx} = \frac{(-\sin x) \cdot \log (\sin x) + \cos x \cdot \frac{1}{\sin x} \cdot \cos x}{1}$$

$$\Rightarrow \frac{dv}{dx} = (\sin x)^{\cos x} \cdot \left[ -\sin x \cdot \log \sin x + \cos x \cdot \cot x \right]$$

$$\frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$$

$$\frac{dy}{dx} = x^{\sin x} \cdot \left( \cos x \cdot \log x + \frac{\sin x}{x} \right)$$

$$+ (\sin x)^{\cos x} \cdot \left[ -\sin x \cdot \log \sin x + \cos x \cdot \cot x \right]$$



Exercise 5.5 Chapter 5

[Q.10] Differentiate with respect to  $x$ .

$$y = \frac{x^{\cos x}}{x} + \frac{x^2+1}{x^2-1}$$

$$y = u + v$$

$$\frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$$

Var.  $y = \frac{\text{Var.}}{\text{Var.}}$   
 log. Diff.  $\log(m^n) = n \cdot \log m$   
 $\log \rightarrow \text{Diff.}$

$$(u \cdot v)' = u' \cdot v + u \cdot v'$$

$$(u \cdot v \cdot w)' = u' \cdot vw + u \cdot v'w + u \cdot v \cdot w'$$

$$u = x^{\cos x}$$

$$\Rightarrow \log u = \log(x^{\cos x})$$

$$\Rightarrow \log u = \cos x \cdot \log x$$

Diff. w.r.t(x)

$$\Rightarrow \frac{1}{u} \cdot \left( \frac{du}{dx} \right) = 1 \cdot \cos x \cdot \log x + x(-\sin x) \log x + x^{\cos x} \cdot \frac{1}{x}$$

$$\Rightarrow \frac{du}{dx} = x^{\cos x} \left[ \cos x \cdot \log x - x \sin x \cdot \log x + \cos x \right]$$

now  $v = \frac{x^2+1}{x^2-1}$

$$\left( \frac{u}{v} \right)' = \frac{u'v - u \cdot v'}{v^2}$$

$$\Rightarrow \frac{dv}{dx} = \frac{(2x+0) \cdot (x^2-1) - (x^2+1) \cdot (2x)}{(x^2-1)^2}$$

$$\frac{dv}{dx} = \frac{2x^3 - 2x - 2x^3}{(x^2-1)^2}$$

$$\frac{dv}{dx} = \frac{-4x}{(x^2-1)^2}$$



$$\frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$$

$$\frac{dy}{dx} = x^{\cos x} \left[ \cos x \cdot \log x - x \sin x \log x + \cos x \right] - \frac{4x}{(x^2-1)^2}$$

Q.11  $y = (x \cos x)^n + (x \sin x)^{1/n}$

$$y = \underbrace{(x \cos x)^n}_u + \underbrace{(x \sin x)^{1/n}}_v \Rightarrow \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$$

$$u = (x \cos x)^n$$

$$\Rightarrow \log u = \log (x \cos x)^n$$

$$\Rightarrow \log u = \underline{n \cdot \log (x \cos x)}$$

Diff. w.r.t.  $x$

$$\Rightarrow \frac{1}{u} \cdot \frac{du}{dx} = \frac{1 \cdot \log (x \cos x)}{x \cos x} + \cancel{n} \cdot \frac{1}{x \cos x} \cdot [1 \cdot \cos x + x \cdot (-\sin x)]$$

$$\Rightarrow \frac{du}{dx} = (x \cos x)^n \cdot [\log (x \cos x) + 1 - x \cdot \tan x] \quad \text{--- (i)}$$

Now  $V = (x \sin x)^{1/n}$

$$\Rightarrow \log V = \log (x \sin x)^{1/n}$$

$$\Rightarrow \log V = \frac{1}{n} \cdot \log (x \sin x)$$

by Diff. w.r.t.  $(x) \rightarrow$



$$\Rightarrow \frac{1}{v} \cdot \left( \frac{dv}{dn} \right) = \left( -\frac{1}{n^2} \right) \cdot \log(n \sin n) + \frac{1}{n} \cdot \frac{1}{n \sin n} \cdot (1 \cdot \sin n + n \cdot \cos n)$$

$$\Rightarrow \frac{dv}{dn} = (n \sin n)^{\frac{1}{n}} \cdot \left[ -\frac{\log(n \sin n)}{n^2} + \frac{1}{n^2} \cdot (1 + n \cdot \cot n) \right]$$

$$\Rightarrow \frac{dv}{dn} = (n \sin n)^{\frac{1}{n}} \cdot \left[ \frac{-\log(n \sin n) + 1 + n \cot n}{n^2} \right] \quad \text{--- (2)}$$

$$\frac{dy}{dn} = \frac{du}{dn} + \frac{dv}{dn} \quad \text{by eqn (1) \& (2)}$$

$$\Rightarrow \frac{dy}{dn} = (n \cos n)^n \cdot \left[ \log(n \cos n) + 1 - n \tan n \right] + (n \sin n)^{\frac{1}{n}} \cdot \left[ \frac{-\log(n \sin n) + 1 + n \cot n}{n^2} \right]$$



$$\Rightarrow \frac{dy}{dn} = \frac{-\left(x^y \cdot \frac{y}{x} + y^x \cdot \log y\right)}{\left(x^y \cdot \log x + y^x \cdot \frac{x}{y}\right)}$$

$$\Rightarrow \frac{dy}{dn} = - \frac{\left(x^{y-1} \cdot y + y^x \cdot \log y\right)}{\left(x^y \cdot \log x + y^{x-1} \cdot x\right)}$$

$$\left(\frac{y}{x}\right) \cdot \frac{y}{x}$$

$$x^{y-1} \cdot y$$

[Q.13]  $y^x = x^y$

$$\Rightarrow \log(y^x) = \log(x^y)$$

$$\Rightarrow x \cdot \log y = y \cdot \log x$$

by diff. w.r.t.  $(x) \rightarrow$

$$\Rightarrow 1 \cdot \log y + x \cdot \frac{1}{y} \cdot \frac{dy}{dx} = \frac{dy}{dx} \cdot \log x + y \cdot \frac{1}{x}$$

$$\Rightarrow \left(\frac{dy}{dx}\right) \left(\frac{x}{y} - \log x\right) = \left(\frac{y}{x} - \log y\right)$$

$$\Rightarrow \frac{dy}{dx} = \frac{\frac{y}{x} - \log y}{\frac{x}{y} - \log x} = \frac{(y - x \log y)/x}{(x - y \log x)/y}$$

$$\frac{dy}{dx} = \frac{y}{x} \cdot \left(\frac{y - x \log y}{x - y \log x}\right)$$



Q.14

$$(\cos x)^y = (\cos y)^x$$

$$\Rightarrow \log (\cos x)^y = \log (\cos y)^x$$

$$\Rightarrow \frac{y \cdot \log (\cos x)}{\text{diff. w.r.t. } (x)} = \frac{x \cdot \log (\cos y)}{\text{diff. w.r.t. } (x)} \rightarrow$$

$$\Rightarrow \left( \frac{dy}{dx} \right) \cdot \log \cos x + y \cdot \frac{1}{\cos x} \cdot (-\sin x) = 1 \cdot \log (\cos y) + x \cdot \frac{1}{\cos y} \cdot (-\sin y) \cdot \frac{dy}{dx}$$

$$\Rightarrow \left( \frac{dy}{dx} \right) (\log \cos x + x \cdot \tan y) = \log (\cos y) + x \tan x$$

$$\Rightarrow \boxed{\frac{dy}{dx} = \frac{\log \cos y + x \tan x}{\log \cos x + x \tan y}}$$

Q.15

$$xy = e^{(x-y)}$$

$$\Rightarrow \log (xy) = \log (e^{(x-y)})$$

$$\Rightarrow \log x + \log y = (x-y) \cdot (\log_e e)$$

$$\Rightarrow \log x + \log y = x-y$$


Diff. w.r.t. (x)

$$\Rightarrow \frac{1}{x} + \frac{1}{y} \cdot \frac{dy}{dx} = 1 - \frac{dy}{dx}$$

$$\Rightarrow \left( \frac{dy}{dx} \right) \left( \frac{1}{y} + 1 \right) = 1 - \frac{1}{x}$$

$$\frac{dy}{dx} = \frac{\left( 1 - \frac{1}{x} \right)}{\left( \frac{1}{y} + 1 \right)} = \frac{\left( \frac{x-1}{x} \right)}{\left( \frac{1+y}{y} \right)}$$

$$\Rightarrow \boxed{\frac{dy}{dx} = \frac{y(x-1)}{x(1+y)}}$$





Exercise-5.5 Chapter-5

[Q.16] Find the derivative of the function given by  $f(x) = (1+x)(1+x^2)(1+x^4)(1+x^8)$  and hence find  $f'(1)$ .

Sol<sup>n</sup>  $f(x) = (1+x) \cdot (1+x^2) \cdot (1+x^4) \cdot (1+x^8)$

$$\Rightarrow \log(f(x)) = \log[(1+x)(1+x^2)(1+x^4)(1+x^8)]$$

$$\Rightarrow \log f(x) = \log(1+x) + \log(1+x^2) + \log(1+x^4) + \log(1+x^8)$$

by Diff. w.r.t. 'x'

$$\Rightarrow \frac{1}{f(x)} \cdot \left( \frac{d(f(x))}{dx} \right) = \left[ \frac{1 \cdot (0+1)}{1+x} + \frac{1 \cdot (0+2x)}{1+x^2} + \frac{1 \cdot (0+4x^3)}{1+x^4} + \frac{1 \cdot (0+8x^7)}{1+x^8} \right]$$

$f'(x)$

$$\Rightarrow f'(x) = (1+x)(1+x^2)(1+x^4)(1+x^8) \left[ \frac{1}{1+x} + \frac{2x}{1+x^2} + \frac{4x^3}{1+x^4} + \frac{8x^7}{1+x^8} \right]$$

Put  $x=1$

$$f'(1) = (1+1)(1+1)(1+1)(1+1) \left[ \frac{1}{2} + \frac{2}{2} + \frac{4}{2} + \frac{8}{2} \right]$$

$$f'(1) = 8 \times \frac{15}{2} = 120$$



Q.17

let  $y = (x^2 - 5x + 8)(x^3 + 7x + 9)$

(i) Differentiate by using product rule.

$$(u.v)' = u'.v + u.v'$$

$$y = \overset{u}{(x^2 - 5x + 8)} \cdot \overset{v}{(x^3 + 7x + 9)}$$

$$\frac{dy}{dx}$$

diff. w.r.t. x

$$\frac{dy}{dx} = (\underline{2x} - 5) \cdot (x^3 + 7x + 9) + (x^2 - 5x + 8) \cdot (3x^2 + 7)$$

$$= \underline{2x^4} + \underline{14x^2} + \underline{18x} - \underline{5x^3} - \underline{35x} - 45 + \underline{3x^4} + \underline{7x^2} - \underline{15x^3} - \underline{35x} + \underline{24x^2} + 56$$

$$\frac{dy}{dx} = \underline{5x^4 - 20x^3 + 45x^2 - 52x + 11}$$

(ii) by expanding the product to obtain a single polynomial

$$y = (x^2 - 5x + 8)(x^3 + 7x + 9)$$

$$\Rightarrow y = \underline{x^5} + \underline{7x^3} + \underline{9x^2} - \underline{5x^4} - \underline{35x^2} - \underline{45x} + \underline{8x^3} + \underline{56x} + 72$$

$$\Rightarrow y = \underline{x^5 - 5x^4 + 15x^3 - 26x^2 + 11x + 72}$$

$$\frac{dy}{dx} = 5x^4 - 20x^3 + 45x^2 - 52x + 11 + 0$$



(iii) by logarithmic differentiation.

$$y = (x^2 - 5x + 8)(x^3 + 7x + 9)$$

$$\Rightarrow \log y = \log [(x^2 - 5x + 8) \cdot (x^3 + 7x + 9)]$$

$$\log(m \cdot n) = \log m + \log n$$

$$\Rightarrow \log y = \log(x^2 - 5x + 8) + \log(x^3 + 7x + 9)$$

by Diff. w.r.t.  $(x) \rightarrow$

$$\Rightarrow \frac{1}{y} \cdot \left( \frac{dy}{dx} \right) = \left[ \frac{1}{x^2 - 5x + 8} \times (2x - 5) + \frac{1}{x^3 + 7x + 9} \cdot (3x^2 + 7) \right]$$

$$\Rightarrow \frac{dy}{dx} = \frac{(x^2 - 5x + 8) \cdot (2x - 5) + (3x^2 + 7)(x^3 + 7x + 9)}{(x^2 - 5x + 8)(x^3 + 7x + 9)}$$

$$\Rightarrow \frac{dy}{dx} = (2x - 5)(x^3 + 7x + 9) + (3x^2 + 7)(x^2 - 5x + 8)$$

$$\Rightarrow \boxed{\frac{dy}{dx} = 5x^4 - 20x^3 + 45x^2 - 52x + 11}$$

(yes)



Q.18

$$\frac{d}{dn} (u.v.w) = \frac{du}{dn} \cdot v.w + u \cdot \frac{dv}{dn} \cdot w + u.v \cdot \frac{dw}{dn}$$

(i) Prove by repeated application of Product rule.

$$\text{LHS} = \frac{d}{dn} (u.v.w) = \frac{d}{dn} [\underline{u} \cdot (\underline{v.w})]$$

$$= \frac{du}{dn} \cdot (v.w) + u \cdot \frac{d(v.w)}{dn}$$

$$= \frac{du}{dn} \cdot v.w + u \cdot \left( \frac{dv}{dn} \cdot w + v \cdot \frac{dw}{dn} \right)$$

$$= \underbrace{\frac{du}{dn} \cdot v.w} + \underbrace{u \cdot \frac{dv}{dn} \cdot w} + \underbrace{u.v \cdot \frac{dw}{dn}} = \text{RHS.}$$

(ii) Prove by logarithmic differentiation.

Let  $y = u.v.w$

$$\frac{d(u.v.w)}{dn}$$

$$\Rightarrow \log y = \log (u.v.w)$$

$$\Rightarrow \log y = \log u + \log v + \log w$$

~~diff~~ diff. w.r.t  $(x)$

$$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dn} = \frac{1}{u} \cdot \frac{du}{dn} + \frac{1}{v} \cdot \frac{dv}{dn} + \frac{1}{w} \cdot \frac{dw}{dn}$$

$$\Rightarrow \frac{dy}{dn} = (u.v.w) \left( \frac{1}{u} \cdot \frac{du}{dn} + \frac{1}{v} \cdot \frac{dv}{dn} + \frac{1}{w} \cdot \frac{dw}{dn} \right)$$

$$\Rightarrow \frac{d(u.v.w)}{dn} = \frac{du}{dn} \cdot v.w + u \cdot \frac{dv}{dn} \cdot w + u.v \cdot \frac{dw}{dn}$$



## Derivatives of Functions in Parametric Form (प्राचलिक रूप)

### Functions

$y \leftrightarrow x$

Explicit Function

$$y = x^2 + 3x + 1$$

$$y = \sin x$$

$$y = e^{\tan^{-1}(x^2)}$$

Implicit Function

$$xy^3 = y + \sin x$$

$$y + \tan^{-1} x = \sec(\pi y)$$

Parametric Form

$x \leftrightarrow y$   
Parameter  
New Variable  
 $t, x, \theta$  etc.

e.g.  $\begin{cases} y = 2 + 4t \\ x = 7t^2 \end{cases} \rightarrow t = \text{Parameter}$

e.g.  $\begin{cases} y = 2 \sin \theta \\ x = 5 \cos \theta \end{cases} \rightarrow \theta = \text{Parameter}$

How to find  $\frac{dy}{dx}$  if  $x = f(t)$ ,  $y = g(t)$ ?

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)}$$

$x \leftrightarrow y$   
 $t$

Diff. w.r. to 't'

$$y = g(t) \rightarrow \frac{dy}{dt} = g'(t)$$

$$x = f(t) \rightarrow \frac{dx}{dt} = f'(t)$$



e.g. Find  $\frac{dy}{dx}$ , if  $x = a \cos \theta$ ,  $y = a \sin \theta$ .

$a = \text{Constant}$  ( $\text{সংকট}$ )

$\theta \rightarrow \text{Parameter}$  ( $\text{প্যারামিটার}$ )

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{d\theta}\right)}{\left(\frac{dx}{d\theta}\right)} = \frac{a \cos \theta}{-a \sin \theta}$$

$$\boxed{\frac{dy}{dx} = -\cot \theta}$$

$$y = a \sin \theta$$

diff. w.r.t ' $\theta$ '

$$\frac{dy}{d\theta} = a \cos \theta \quad \text{--- (1)}$$

$$x = a \cos \theta$$

diff. w.r.t.  $\theta$

$$\frac{dx}{d\theta} = -a \sin \theta \quad \text{--- (2)}$$

↳ Attached Question.

Find  $\frac{dy}{dx}$  in terms of  $x$  &  $y$ .

$$\frac{dy}{dx} = -\cot \theta = \frac{a \cos \theta}{-a \sin \theta}$$

$$\Rightarrow \frac{dy}{dx} = -\frac{x}{y}$$

$$\Rightarrow \boxed{\frac{dy}{dx} = -\frac{x}{y}}$$

$$\begin{aligned} x &= a \cos \theta \\ y &= a \sin \theta \end{aligned}$$

Given  
Substitution



## Alternate way of solving attached Question

$$\begin{cases} x = a \cos \theta \\ y = a \sin \theta \end{cases}$$

find  $\frac{dy}{dx}$  in terms of  $x$  &  $y$ .

$$\boxed{\cos^2 \theta + \sin^2 \theta = 1}$$

$$x^2 = a^2 \cos^2 \theta \quad \text{--- (1)}$$

$$+ y^2 = a^2 \sin^2 \theta \quad \text{--- (2)}$$

$$x^2 + y^2 = a^2 (\cos^2 \theta + \sin^2 \theta)$$

$$\Rightarrow \boxed{x^2 + y^2 = a^2}$$

by Diff. w.r.t 'x'

$$\Rightarrow 2x + 2y \left( \frac{dy}{dx} \right) = 0$$

$$\Rightarrow \frac{dy}{dx} = \frac{-2x}{2y} = \frac{-x}{y}$$


Note,

$$\boxed{x^2 + y^2 = a^2}$$

Circle in Cartesian Form

Centre  $(0, 0)$

Radius  $a$



$$\begin{cases} x = a \cos \theta \\ y = a \sin \theta \end{cases}$$

Circle in  
Parametric Form



Exercise 5.6 Chapter-5

Parametric form  $\star \left( \frac{dy}{dx} = \frac{\left( \frac{dy}{dt} \right)}{\left( \frac{dx}{dt} \right)} \right)$

$x \swarrow \searrow y$   
 $\downarrow \downarrow$   
 Parameter  $(x, t, \theta \dots)$   
 $(t)$

[Q.1]  $x = 2at^2, y = at^4$

Diff. w.r.to 't'  $\downarrow$

$$\frac{dx}{dt} = 2a(2t) = 4at, \quad \frac{dy}{dt} = 4at^3$$

$$\frac{dy}{dx} = \frac{\left( \frac{dy}{dt} \right)}{\left( \frac{dx}{dt} \right)} = \frac{4at^3}{4at} = t^2$$

[Q.2]  $x = a \cos \theta, y = b \cos \theta$

$$\frac{dy}{dx} = \frac{\left( \frac{dy}{d\theta} \right)}{\left( \frac{dx}{d\theta} \right)} = \frac{\frac{d(b \cos \theta)}{d\theta}}{\frac{d(a \cos \theta)}{d\theta}} = \frac{b(-\sin \theta)}{a(-\sin \theta)} = \frac{b}{a}$$

[Q.3]  $x = \sin t, y = \cos 2t$

$$\frac{dy}{dx} = \frac{\left( \frac{dy}{dt} \right)}{\left( \frac{dx}{dt} \right)} = \frac{\left( \frac{d(\cos 2t)}{dt} \right)}{\left( \frac{d(\sin t)}{dt} \right)} = \frac{-\sin 2t \cdot (2)}{\cos t}$$

$$= \frac{-2 \sin t \cdot \cos t \times 2}{\cos t}$$

$$= -4 \sin t$$



$$\boxed{Q.4} \quad x = 4t, \quad y = \frac{4}{t}$$

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{\frac{d\left(\frac{4}{t}\right)}{dt}}{\frac{d(4t)}{dt}} = \frac{\cancel{4}\left(-\frac{1}{t^2}\right)}{\cancel{4} \times 1} = -\frac{1}{t^2}$$

$$\boxed{Q.5} \quad x = \cos\theta - \cos 2\theta, \quad y = \sin\theta - \sin 2\theta$$

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{d\theta}\right)}{\left(\frac{dx}{d\theta}\right)} = \frac{\frac{d(\sin\theta - \sin 2\theta)}{d\theta}}{\frac{d(\cos\theta - \cos 2\theta)}{d\theta}} = \frac{\cos\theta - 2\cos 2\theta}{-\sin\theta + \sin 2\theta \cdot 2}$$

✓

$$\boxed{Q.6} \quad x = a(\theta - \sin\theta), \quad y = a(1 + \cos\theta)$$

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{d\theta}\right)}{\left(\frac{dx}{d\theta}\right)} = \frac{\frac{d[a(1 + \cos\theta)]}{d\theta}}{\frac{d[a(\theta - \sin\theta)]}{d\theta}} = \frac{\cancel{a}(0 - \sin\theta)}{\cancel{a}(1 - \cos\theta)}$$

$$\frac{dy}{dx} = -\left(\frac{\sin\theta}{1 - \cos\theta}\right) = -\left(\frac{\cancel{2}\sin\frac{\theta}{2} \cdot \cos\frac{\theta}{2}}{\cancel{2}\sin^2\frac{\theta}{2}}\right)$$

$$\cos\theta = 1 - 2\sin^2\frac{\theta}{2}$$

$$= -\frac{\cos\frac{\theta}{2}}{\sin\frac{\theta}{2}} = -\cot\frac{\theta}{2}$$



Q. 7

$$x = \frac{\sin^3 t}{\sqrt{\cos 2t}}$$

$$y = \frac{\cos^3 t}{\sqrt{\cos 2t}}$$

$$\frac{dy}{dx} = \frac{(dy/dt)}{(dx/dt)} =$$

$$y = \frac{\cos^3 t}{\sqrt{\cos 2t}}$$

$$\left(\frac{u}{v}\right)' = \frac{u' \cdot v - u \cdot v'}{v^2}$$

by diff. w.r. to 't'

$$\frac{dy}{dt} = \frac{3\cos^2 t \cdot (-\sin t) \cdot \sqrt{\cos 2t} - \cos^3 t \cdot \frac{1}{\sqrt{\cos 2t}} \cdot (-\sin 2t)}{(\sqrt{\cos 2t})^2}$$

$$\frac{dy}{dt} = \frac{\left( \frac{-3\cos^2 t \cdot \sin t \cdot \cos 2t + \cos^3 t \cdot \sin 2t}{\sqrt{\cos 2t}} \right)}{\cos 2t}$$

$$\frac{dy}{dt} = \frac{-3\cos^2 t \cdot \sin t \cdot \cos 2t + \cos^3 t \cdot \sin 2t}{\cos 2t \cdot \sqrt{\cos 2t}} \quad \text{--- (1)}$$



$$x = \frac{\sin^3 t}{\sqrt{\cos 2t}}$$

$$\left(\frac{u}{v}\right)' = \frac{u'v - u.v'}{v^2}$$

by diff. w.r. to 't'

$$\frac{dx}{dt} = \frac{3\sin^2 t \cdot \cos t \cdot \sqrt{\cos 2t} - \sin^3 t \cdot \frac{1}{\sqrt{\cos 2t}} \cdot (-\sin 2t)}{\cos 2t}$$

$$\frac{dx}{dt} = \frac{3\sin^2 t \cdot \cos t \cdot \cos 2t + \sin^3 t \cdot \sin 2t}{\cos 2t \cdot \sqrt{\cos 2t}}$$

$$\Rightarrow \frac{dx}{dt} = \frac{3\sin^2 t \cdot \cos t \cdot \cos 2t + \sin^3 t \cdot \sin 2t}{\cos 2t \cdot \sqrt{\cos 2t}} \quad \text{--- (2)}$$

$$\frac{dy}{dx} = \frac{(dy/dt)}{(dx/dt)}$$

(by putting the values of  $\frac{dy}{dt}$  &  $\frac{dx}{dt}$  in ~~eq~~ from eqn ① & ②)

$$\Rightarrow \frac{dy}{dx} = \frac{-3\cos^2 t \cdot \sin t \cdot \boxed{\cos 2t} + \cos^3 t \cdot \boxed{\sin 2t}}{\cos 2t \cdot \sqrt{\cos 2t}}$$

$$\frac{3\sin^2 t \cdot \cos t \cdot \boxed{\cos 2t} + \sin^3 t \cdot \boxed{\sin 2t}}{\cos 2t \cdot \sqrt{\cos 2t}}$$

put  $\boxed{\sin 2t = 2 \sin t \cdot \cos t}$



$$\Rightarrow \frac{dy}{dn} = \frac{-3\cos^2 t \cdot \sin t \cdot \boxed{\cos 2t} + \cos^3 t \cdot (2\sin t \cdot \cos t)}{3\sin^2 t \cdot \cos t \cdot \boxed{\cos 2t} + \sin^3 t \cdot (2\sin t \cdot \cos t)}$$

$$\Rightarrow \frac{dy}{dn} = \frac{\cancel{\sin t} \cdot \cos^2 t \cdot [-3 \cdot \cancel{\cos 2t} + 2\cos^2 t]}{\sin^2 t \cdot \cancel{\cos t} [3\cancel{\cos 2t} + 2\sin^2 t]}$$

$$\Rightarrow \frac{dy}{dn} = \frac{\cos t [-3(2\cos^2 t - 1) + 2\cos^2 t]}{\sin t [3(1 - 2\sin^2 t) + 2\sin^2 t]}$$

$$\Rightarrow \frac{dy}{dn} = \frac{\cos t \cdot [-4\cos^2 t + 3]}{\sin t \cdot [3 - 4\sin^2 t]}$$

$$\Rightarrow \frac{dy}{dn} = \frac{-4\cos^3 t + 3\cos t}{3\sin t - 4\sin^3 t} = \frac{-\cos 3t}{\sin 3t}$$

$$\boxed{\frac{dy}{dn} = -\cot 3t}$$



Q.8

$$x = a \left( \cos t + \log \tan \frac{t}{2} \right), \quad y = a \sin t$$

$$\frac{dy}{dx} = \frac{\left( \frac{dy}{dt} \right)}{\left( \frac{dx}{dt} \right)} = \frac{\frac{d(a \sin t)}{dt}}{\frac{d\left(a \left( \cos t + \log \tan \frac{t}{2} \right) \right)}{dt}}$$

$$= \frac{a \cos t}{a \left( -\sin t + \frac{1}{\tan \frac{t}{2}} \cdot \sec^2 \frac{t}{2} \cdot \frac{1}{2} \right)}$$

$$= \frac{\cos t}{\left( -\sin t + \frac{\cos t}{2 \sin \frac{t}{2}} \cdot \frac{1}{\cos^2 \frac{t}{2}} \right)} = \frac{\cos t}{\left( -\sin t + \frac{1}{\sin t} \right)}$$

$$= \frac{\cos t}{\left( \frac{-\sin^2 t + 1}{\sin t} \right)} = \frac{\cancel{\cos t} \cdot \sin t}{\cos^2 t}$$

$$= \tan t$$

$$\frac{dy}{dx} = \tan t$$



**[Q.9]**  $x = a \sec \theta$  ,  $y = b \tan \theta$

$$\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{\left( \frac{d(b \tan \theta)}{d\theta} \right)}{\left( \frac{d(a \sec \theta)}{d\theta} \right)} = \frac{b \sec^2 \theta}{a \sec \theta \cdot \tan \theta}$$

$$\frac{dy}{dx} = \frac{b}{a} \frac{\left( \frac{1}{\cancel{\cos \theta}} \right)}{\left( \frac{\sin \theta}{\cancel{\cos \theta}} \right)} = \frac{b}{a} \frac{1}{\sin \theta} = \frac{b}{a} \operatorname{cosec} \theta$$

**[Q.10]**  $x = a(\cos \theta + \theta \sin \theta)$  ,  $y = a(\sin \theta - \theta \cos \theta)$

$$\frac{dy}{dx} = \frac{\left( \frac{dy}{d\theta} \right)}{\left( \frac{dx}{d\theta} \right)} = \frac{\frac{d(a(\sin \theta - \theta \cdot \cos \theta))}{d\theta}}{\frac{d(a(\cos \theta + \theta \cdot \sin \theta))}{d\theta}}$$

$$= \frac{a[\cancel{\cos \theta} - 1 \cdot \cos \theta + \theta \cdot \sin \theta]}{a[-\sin \theta + 1 \cdot \sin \theta + \theta \cdot \cos \theta]}$$

$$= \frac{\theta \cdot \sin \theta}{\theta \cdot \cos \theta} = \tan \theta \quad \checkmark$$



**[Q.11]** If  $x = \sqrt{a^{\sin^{-1}t}}$ ,  $y = \sqrt{a^{\cos^{-1}t}}$ , then show that

$$\boxed{x = a^{\frac{\sin^{-1}t}{2}}}, \quad \boxed{y = a^{\frac{\cos^{-1}t}{2}}}, \quad \boxed{\frac{dy}{dx} = -\frac{y}{x}}$$

I. method

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{\frac{d\left(a^{\frac{\cos^{-1}t}{2}}\right)}{dt}}{\frac{d\left(a^{\frac{\sin^{-1}t}{2}}\right)}{dt}} = \frac{a^{\frac{\cos^{-1}t}{2}} \cdot \log a \cdot \frac{1}{2} \cdot \frac{-1}{\sqrt{1-t^2}}}{a^{\frac{\sin^{-1}t}{2}} \cdot \log a \cdot \frac{1}{2} \cdot \frac{1}{\sqrt{1-t^2}}}$$

$$\boxed{\frac{d(a^x)}{dx} = a^x \cdot \log a}$$

$$= -\frac{y}{x}$$

II. method

$x = a^{\frac{\sin^{-1}t}{2}}$ ,  $y = a^{\frac{\cos^{-1}t}{2}}$  (logarithmic Diff.)

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)}$$

$$= \frac{-y}{\sqrt{1-t^2}} \cdot \frac{\log a}{2}$$

$$\frac{x}{\sqrt{1-t^2}} \cdot \frac{\log a}{2}$$

$$= -\frac{y}{x}$$

$$\begin{aligned} x &= a^{\frac{\sin^{-1}t}{2}} \\ \Rightarrow \log x &= \frac{\sin^{-1}t}{2} \cdot \log a \\ &\text{diff. w.r.to } t \\ \Rightarrow \frac{1}{x} \cdot \frac{dx}{dt} &= \frac{1}{\sqrt{1-t^2}} \cdot \frac{\log a}{2} \end{aligned}$$

$$\Rightarrow \boxed{\frac{dx}{dt} = \frac{x}{\sqrt{1-t^2}} \cdot \frac{\log a}{2}}$$

Similarly

$$\frac{dy}{dt} = \frac{-y}{\sqrt{1-t^2}} \cdot \frac{\log a}{2}$$



III - method

$$x = a^{\frac{\sin^{-1}t}{2}} \quad (1), \quad y = a^{\frac{\cos^{-1}t}{2}} \quad (2)$$

To prove,

$$\frac{dy}{dx} = -\frac{y}{x}$$

Property

$$\sin^{-1}t + \cos^{-1}t = \frac{\pi}{2}$$

By eq<sup>n</sup> (1)  $\times$  eq<sup>n</sup> (2)  $\rightarrow$

$$xy = a^{\frac{\sin^{-1}t}{2}} \cdot a^{\frac{\cos^{-1}t}{2}}$$

$$\Rightarrow xy = a^{\frac{1}{2}(\sin^{-1}t + \cos^{-1}t)} \rightarrow \left(\frac{\pi}{2}\right)$$

$$\Rightarrow \boxed{xy = a^{\frac{\pi}{4}}}$$

by diff. w.r.t. 'x'

$$\Rightarrow 1 \cdot y + x \cdot \left(\frac{dy}{dx}\right) = 0$$

$$\Rightarrow \boxed{\frac{dy}{dx} = -\frac{y}{x}} \quad \checkmark$$



## Second Order Derivative (द्वितीय कोटि अवकलन)

$y = f(x) \rightarrow$  Function

$$D^1 y = y' = y_1 = \frac{dy}{dx} = \frac{d(f(x))}{dx} = f'(x) \rightarrow \text{First order Derivative}$$

$$D^2 y = y'' = y_2 = \frac{d^2 y}{dx^2} = \frac{d^2(f(x))}{dx^2} = \frac{d\left(\frac{d(f(x))}{dx}\right)}{dx} = f''(x)$$

$\rightarrow$  Second order Derivative

$y = f(x) \rightarrow$  Function

$$\frac{dy}{dx} = y' = f'(x) \rightarrow \text{slope of function}$$

$$\frac{d^2 y}{dx^2} = y'' = f''(x) \rightarrow \text{Concavity of the function.}$$

Concave  $\downarrow \ominus$   
Convex  $\downarrow \oplus$

| <u>Distance</u><br>(s) | <u>Velocity</u><br>(v) | <u>Acceleration</u><br>(a)                                                         |
|------------------------|------------------------|------------------------------------------------------------------------------------|
|                        | $v = \frac{ds}{dt}$    | $a = \frac{dv}{dt} = \frac{d\left(\frac{ds}{dt}\right)}{dt} = \frac{d^2(s)}{dt^2}$ |

e.g. If  $y = \sin^{-1} x$ , then prove that  $\boxed{(1-x^2)y'' - x.y' = 0}$

$$y = \sin^{-1} x$$

Diff. w.r.t. (x)  $\rightarrow$

$$y' = \frac{dy}{dx} = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d^2 y}{dx^2} \quad \frac{dy}{dx}$$



$$y = \sin^{-1} x$$

$$y' = \frac{dy}{dx} = \frac{1}{\sqrt{1-x^2}}$$

Again diff. w.r.t.  $(x) \rightarrow$

$$y'' = \frac{d^2 y}{dx^2} = \frac{d\left(\frac{1}{\sqrt{1-x^2}}\right)}{dx} = \frac{0 \cdot \sqrt{1-x^2} - 1 \cdot \left(\frac{1}{\sqrt{1-x^2}}\right) \cdot (-2x)}{(1-x^2)^2}$$

$$y'' = + \frac{x}{(1-x^2)\sqrt{1-x^2}}$$

$$\Rightarrow (1-x^2) \cdot y'' = \frac{x}{\sqrt{1-x^2}}$$

$$\Rightarrow (1-x^2) \cdot y'' = x \cdot y'$$

$$\Rightarrow \boxed{(1-x^2) \cdot y'' - x \cdot y' = 0} \quad \text{H.P.}$$

Fast.

$$y = \sin^{-1} x$$

Diff.

$$\Rightarrow y' = \frac{1}{\sqrt{1-x^2}}$$

$$\Rightarrow (\sqrt{1-x^2}) \cdot y' = 1$$

Diff.

Square

$$\Rightarrow \frac{(1-x^2) \cdot (y')^2}{(u \cdot v)'} = 1$$

Diff.

$$\frac{(-2x) \cdot (y')^2 + (1-x^2) \cdot 2(y') \cdot (y'')}{\text{}} = 0$$

$$\Rightarrow 2y'(-xy' + (1-x^2) \cdot y'') = 0$$

$$\Rightarrow \boxed{(1-x^2) \cdot y'' - xy' = 0}$$



e.g. If  $x = a \cos \theta$ ,  $y = b \sin \theta$ , Find  $\frac{d^2y}{dx^2}$ .

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{d\theta}\right)}{\left(\frac{dx}{d\theta}\right)}$$

$$\frac{dy}{dx} = \frac{b \cos \theta}{-a \sin \theta}$$

$$y = b \sin \theta$$

$$\frac{dy}{d\theta} = b \cos \theta \quad \checkmark$$

$$x = a \cos \theta$$

$$\frac{dx}{d\theta} = -a \sin \theta$$

$$\frac{dy}{dx} = -\frac{b}{a} \cdot \cot \theta$$

by diff. w.r.t.  $x$   $\rightarrow$

$$\Rightarrow \frac{d^2y}{dx^2} = -\frac{b}{a} \cdot (-\operatorname{cosec}^2 \theta) \cdot \frac{d\theta}{dx}$$

$$\Rightarrow \frac{d^2y}{dx^2} = \frac{b}{a} (\operatorname{cosec}^2 \theta) \cdot \left(\frac{1}{-a \sin \theta}\right)$$


$$\frac{d^2y}{dx^2} = -\frac{b}{a^2} \cdot \operatorname{cosec}^3 \theta$$

$$\frac{d^2y}{dx^2} = ?$$

$$x, y, \theta$$

$$\frac{dx}{d\theta} = -a \sin \theta$$



Exercise-5.7 Chapter-5Find the second order derivatives.**Q.1**

$$\text{let } y = x^2 + 3x + 2$$

$$\Rightarrow \frac{dy}{dx} = 2x + 3$$

$$\Rightarrow \frac{d^2y}{dx^2} = 2$$

**Q.2**

$$y = x^{20}$$

$$\Rightarrow \frac{dy}{dx} = 20 \cdot x^{19}$$

$$\Rightarrow \frac{d^2y}{dx^2} = 20 \cdot 19 \cdot x^{18} = 380x^{18}$$

**Q.3**

$$y = x \cos x$$

(u.v)'

$$\Rightarrow \frac{dy}{dx} = 1 \cdot \cos x + x(-\sin x) = \cos x - \frac{x \sin x}{(u.v)'}$$

$$\Rightarrow \frac{d^2y}{dx^2} = -\sin x - (1 \cdot \sin x + x \cdot \cos x)$$

$$\Rightarrow \frac{d^2y}{dx^2} = -2\sin x - x \cos x$$

**Q.4**  $y = \log x$ 

$$\Rightarrow \frac{dy}{dx} = \frac{1}{x}$$

$$\Rightarrow \frac{d^2y}{dx^2} = -\frac{1}{x^2}$$

$$\frac{1}{x} = x^{-1} \rightarrow x^n \rightarrow n \cdot x^{n-1}$$



$$\boxed{\text{Q.5}} \quad y = \underbrace{x^3 \cdot \log x}_{(u \cdot v)'} \quad \underline{(u \cdot v)'}$$

$$\Rightarrow \frac{dy}{dx} = 3x^2 \cdot \log x + x^3 \cdot \frac{1}{x}$$

$$\Rightarrow \frac{dy}{dx} = \underline{3x^2 \cdot \log x} + x^2$$

$$\hookrightarrow \frac{d^2y}{dx^2} = \underline{6x \cdot \log x + 3x^2 \cdot \frac{1}{x}} + 2x$$

$$\hookrightarrow = 6x \cdot \log x + 5x \quad \checkmark$$

$$\boxed{\text{Q.6}} \quad y = \underbrace{e^x \cdot \sin 5x}_{(u \cdot v)'} \quad \frac{d(e^x)}{dx} = e^x \quad \checkmark$$

$$\frac{dy}{dx} = e^x \cdot \sin 5x + e^x \cdot \cos 5x \cdot 5$$

$$\Rightarrow \frac{dy}{dx} = e^x \cdot \sin 5x + 5e^x \cdot \cos 5x$$

$$\Rightarrow \frac{dy}{dx} = \underline{e^x (\sin 5x + 5 \cos 5x)}$$

$$\hookrightarrow \frac{d^2y}{dx^2} = \underline{e^x (\sin 5x + 5 \cos 5x)} + \underline{e^x (5 \cos 5x + 5 \times 5 \times (-\sin 5x))}$$

$$\frac{d^2y}{dx^2} = e^x (-24 \sin 5x + 10 \cos 5x)$$



$$\boxed{\text{Q.7}} \quad y = e^{6x} \cdot \cos 3x$$

$$\frac{dy}{dx} = e^{6x} \cdot 6 \cdot \cos 3x + e^{6x} \cdot (-3 \sin 3x)$$

$$\frac{dy}{dx} = 3 e^{6x} \cdot (2 \cos 3x - \sin 3x)$$

$$\begin{aligned} \Rightarrow \frac{d^2y}{dx^2} &= 3 \cdot e^{6x} \cdot 6 (2 \cos 3x - \sin 3x) \\ &\quad + 3(e^{6x}) \cdot (-2 \sin 3x \cdot 3 - \cos 3x \cdot 3) \end{aligned}$$

$$\begin{aligned} \frac{d^2y}{dx^2} &= 3 e^{6x} [9 \cos 3x - 12 \sin 3x] \\ &= 9 e^{6x} (3 \cos 3x - 4 \sin 3x) \checkmark \end{aligned}$$

$$\boxed{\text{Q.8}} \quad y = \tan^{-1} x$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{(1+x^2)}$$

$$\frac{d\left(\frac{1}{x}\right)}{dx} = -\frac{1}{x^2}$$

$$\Rightarrow \frac{d^2y}{dx^2} = -\frac{1}{(1+x^2)^2} \cdot x(0+2x) = \frac{-2x}{(1+x^2)^2}$$



[Q.9]  $y = \log(\log x)$  (Chain Rule)

$$\Rightarrow \frac{dy}{dx} = \frac{1}{\log x} \cdot \frac{1}{x} = \frac{1}{\underbrace{x \cdot \log x}_{u \cdot v}}$$

$$\hookrightarrow \frac{d^2y}{dx^2} = -\frac{1}{(x \log x)^2} \cdot \left( 1 \cdot \log x + x \cdot \frac{1}{x} \right)$$

$$\Rightarrow \frac{d^2y}{dx^2} = -\frac{(\log x + 1)}{(x \log x)^2}$$

$$\frac{d(\log x)}{dx} = \frac{1}{x}$$

$$\frac{d\left(\frac{1}{x}\right)}{dx} = -\frac{1}{x^2}$$

[Q.10]  $y = \sin(\log x)$  (Chain)

$$\Rightarrow \frac{dy}{dx} = \underbrace{\cos(\log x)}_{(u \cdot v)'} \cdot \frac{1}{x}$$

$$\hookrightarrow \Rightarrow \frac{d^2y}{dx^2} = \underbrace{-\sin(\log x) \cdot \frac{1}{x} \cdot \frac{1}{x}}_{\text{}} + \cos(\log x) \cdot \left(-\frac{1}{x^2}\right)$$

$$\Rightarrow \frac{d^2y}{dx^2} = -\left(\frac{\sin(\log x) + \cos(\log x)}{x^2}\right)$$

[Q.11] If  $y = 5 \cos x - 3 \sin x$ , Prove that  $\frac{d^2y}{dx^2} + y = 0$

$$y = 5 \cos x - 3 \sin x$$

$$\frac{dy}{dx} = -5 \sin x - 3 \cos x$$

$$\Rightarrow \frac{d^2y}{dx^2} = -5 \cos x + 3 \sin x = -\left(\underline{5 \cos x - 3 \sin x}\right)$$

$$\Rightarrow \frac{d^2y}{dx^2} = -y$$

$$\Rightarrow \boxed{\frac{d^2y}{dx^2} + y = 0}$$



Q.12 \* If  $y = \cos^{-1}x$ , Find  $\frac{d^2y}{dx^2}$  in terms of  $y$  alone.

Ans.  $y = \cos^{-1}x$

$$\Rightarrow \cos y = x$$

Diff. w.r.t. 'x'

$$\Rightarrow -\sin y \left( \frac{dy}{dx} \right) = 1$$

$$\Rightarrow \frac{dy}{dx} = \frac{-1}{\sin y} \quad \text{--- (1)}$$

Diff. w.r.t. (x)

$$\Rightarrow + \frac{d^2y}{dx^2} = - \left( \frac{-1}{\sin^2 y} \right) \cdot (\cos y) \cdot \frac{dy}{dx}$$

$$\Rightarrow \frac{d^2y}{dx^2} = \frac{\cos y}{\sin^2 y} \cdot \left( \frac{-1}{\sin y} \right)$$

$$\boxed{\frac{d^2y}{dx^2} = -\cot y \cdot \operatorname{cosec}^2 y}$$



Exercise 5.7 chapter (5)

Q.13

If  $y = 3 \cos(\log x) + 4 \sin(\log x)$ , show that

$$\underbrace{x^2 y_2 + x y_1 + y = 0.}_{\text{}} \quad \left( y_2 = \frac{d^2 y}{dx^2}, \quad y_1 = \frac{dy}{dx} \right)$$

Ans

$$y = 3 \cos(\log x) + 4 \sin(\log x)$$

Diff. w.r.t.  $x$  —

$$\Rightarrow \left( \frac{dy}{dx} \right) = \frac{-3 \sin(\log x)}{x} + \frac{4 \cos(\log x)}{x}$$

$\downarrow$   
 $y_1$

$$\Rightarrow x \cdot y_1 = -3 \sin(\log x) + 4 \cos(\log x)$$

Diff. w.r.t.  $(x)$ 

$$\Rightarrow 1 \cdot y_1 + x \cdot y_2 = \frac{-3 \cos(\log x)}{x} - \frac{4 \sin(\log x)}{x}$$

$$\Rightarrow x y_1 + \underbrace{(x^2 y_2)} = - (3 \cos \log x + 4 \sin \log x)$$

$$\Rightarrow x^2 y_2 + x y_1 = -(y)$$

$$\Rightarrow \boxed{x^2 y_2 + x y_1 + y = 0} \quad \checkmark$$



**Q.14** If  $y = Ae^{mx} + Be^{nx}$ , show that

$$\frac{d^2y}{dx^2} - (m+n)\frac{dy}{dx} + mny = 0$$

Ans.

$$y = Ae^{mx} + Be^{nx}$$

$$\frac{dy}{dx} = mAe^{mx} + nBe^{nx}$$

$$\frac{d^2y}{dx^2} = m^2Ae^{mx} + n^2Be^{nx}$$

$$\text{LHS} = \left(\frac{d^2y}{dx^2}\right) - (m+n)\left(\frac{dy}{dx}\right) + mn(y)$$

$$= m^2Ae^{mx} + n^2Be^{nx} - (m+n)(mAe^{mx} + nBe^{nx}) + mn(Ae^{mx} + Be^{nx})$$

$$= \cancel{m^2Ae^{mx}} + \cancel{n^2Be^{nx}} - \cancel{m^2Ae^{mx}} - \cancel{mnBe^{nx}} - \cancel{mnAe^{mx}} - \cancel{n^2Be^{nx}} + \cancel{mnAe^{mx}} + \cancel{mnBe^{nx}}$$

$$= 0 = \text{RHS.}$$

**Q.15** If  $y = \underline{500 \cdot e^{7x} + 600 \cdot e^{-7x}}$ , show that  $\boxed{\frac{d^2y}{dx^2} = 49y}$

$$\frac{dy}{dx} = \underline{500 \cdot e^{7x} \cdot 7 + 600 \cdot e^{-7x} \cdot (-7)}$$

✓✓

$$\frac{d^2y}{dx^2} = 500 \cdot e^{7x} \cdot (7)^2 + 600 \cdot e^{-7x} \cdot (-7)^2$$

$$\frac{d^2y}{dx^2} = \underline{49} \left( \underline{500e^{7x} + 600e^{-7x}} \right) = 49 \cdot \underset{\textcircled{y}}{y} \quad \checkmark$$



Q.16

If  $e^y(x+1)=1$ , show that  $\frac{d^2y}{dx^2} = \left(\frac{dy}{dx}\right)^2$ .

Sol<sup>n</sup>:

$$e^y(x+1)=1$$

$$(u.v)' = u.v' + u.v'$$

Diff. w.r.t.  $(x) \rightarrow$

$$\Rightarrow \underbrace{e^y}_{(u)} \cdot \underbrace{\frac{dy}{dx}}_{(v)} \cdot (x+1) + \underbrace{e^y}_{(u)} \cdot (1+0) = 0$$

$$\Rightarrow e^y \left( \frac{dy}{dx} \cdot (x+1) + 1 \right) = 0$$

$$\Rightarrow \frac{dy}{dx} (x+1) + 1 = 0 \Rightarrow \boxed{\frac{dy}{dx} = -\frac{1}{(x+1)}}$$

By Diff. w.r.t.  $(x) \rightarrow$

$$\frac{-dy}{dx} = \frac{1}{x+1}$$

$$\frac{d^2y}{dx^2} = -\left(\frac{-1}{(x+1)^2}\right) \Rightarrow \frac{d^2y}{dx^2} = \left(\frac{1}{x+1}\right)^2$$

$$\Rightarrow \boxed{\frac{d^2y}{dx^2} = \left(\frac{-dy}{dx}\right)^2 = \left(\frac{dy}{dx}\right)^2}$$



Q.17. If  $y = (\tan^{-1} x)^2$ , show that

$$(x^2+1)^2 \cdot y_2 + 2x(x^2+1) \cdot y_1 = 2$$

Solution.

$$y = (\tan^{-1} x)^2$$

$$\Rightarrow \frac{dy}{dx} = y_1 = \frac{2(\tan^{-1} x)}{1+x^2}$$

$$\Rightarrow (x^2+1) \cdot y_1 = 2 \tan^{-1} x$$

Diff. w.r.t. (x)  $\rightarrow$


$$\Rightarrow 2x \cdot y_1 + (x^2+1) \cdot y_2 = \frac{2}{1+x^2}$$

$$\Rightarrow \boxed{2x(1+x^2) \cdot y_1 + (x^2+1)^2 \cdot y_2 = 2}$$





Today —  $\left\{ \begin{array}{l} \text{Rolle's Theorem} \\ \text{Mean Value Theorem (मध्यमान प्रमेय)} \end{array} \right.$




Rolle's Theorem:

~~Let  $f: [a, b] \rightarrow \mathbb{R}$~~

Let  $f: [a, b] \rightarrow \mathbb{R}$  be continuous on  $[a, b]$  and differentiable on  $(a, b)$ , such that  $f(a) = f(b)$ , where  $a, b$  are some real numbers. Then there exists some  $c$  in  $(a, b)$  such that  $f'(c) = 0$ .

$(f'(c)) \rightarrow \text{slope at } x=c$



$(\text{Slope} = 0) \rightarrow \text{graph} \rightarrow \text{Horizontal}$

Rolle's

$f: [a, b] \rightarrow \mathbb{R}$

- ① continuous
- ② Differentiable
- ③  $f(a) = f(b)$

④  $f'(c) = 0, c \in (a, b)$



e.g. Verify Rolle's Theorem for the function  $y = x^2 + 2$ ,  
 $x \in [-2, 2]$

Ans.

$$f(x) = x^2 + 2$$


We know that  $f(x) = x^2 + 2$  is continuous  
in  $[-2, 2]$  & differentiable in  $(-2, 2)$ :

$$\begin{aligned} f(a) &= f(-2) = (-2)^2 + 2 = 6 \\ f(b) &= f(2) = (2)^2 + 2 = 6 \end{aligned} \quad \Rightarrow \quad \underline{f(a) = f(b)}$$

$$f(x) = x^2 + 2 \rightarrow \boxed{f'(x) = 2x}$$

$$f'(c) = 2c = 0$$

$$\Rightarrow \boxed{c = 0} \in [-2, 2]$$




$\therefore$  Rolle's Theorem  
is verified.

Mean Value Theorem (MVT) (मध्यमान प्रमेय)

Let  $f: [a, b] \rightarrow \mathbb{R}$  be a continuous function on  $[a, b]$   
and differentiable on  $(a, b)$ .

Then there exists some  $c \in (a, b)$  such that


$$\underbrace{f'(c)} = \underbrace{\frac{f(b) - f(a)}{b - a}}$$



$$m_{AB} = \frac{f(b) - f(a)}{b - a}$$

$$\begin{aligned} \text{Slope of } f' \text{ at } c \\ = \underline{f'(c)} \end{aligned}$$





At some instant

$$\underline{f'(c)} = \underline{30} \text{ speed} = \frac{f(b) - f(a)}{b - a}$$

e.g. Verify mean value theorem for the function  
 $f(x) = x^2$  in the interval  $[2, 4]$

MVT  $\rightarrow$  Continuous  $\checkmark$  in  $[2, 4]$

Differentiable  $\checkmark$  in  $(2, 4) \xrightarrow{a} b$

$$f'(c) = \frac{f(b) - f(a)}{b - a}, \quad \underline{c \in (2, 4)} ??$$

$$f(x) = x^2$$

$$f'(x) = 2x$$

$$\underline{f'(c) = 2c}$$

$$\frac{f(4) - f(2)}{4 - 2} = \frac{16 - 4}{4 - 2} = \frac{12}{2} = 6$$

$$2c = 6$$

$$c = 3 \in (2, 4) \checkmark$$

$\therefore$  MVT  $\checkmark$

Comparison

$$f: [a, b] \rightarrow \mathbb{R}$$

$$\underline{c \in (a, b)}$$

Rolle's Theorem

- ① continuous
- ② differentiable
- ③  $f(a) = f(b)$
- ④  $f'(c) = 0$

MVT

- ① continuous
- ② differentiable
- ③  $f'(c) = \frac{f(b) - f(a)}{b - a}$




Exercise-5.8 Chapter 5

## Rolle's Theorem

$$f: [a, b] \rightarrow \mathbb{R}$$

- ① Continuous in  $[a, b]$
- ② differentiable in  $(a, b)$
- ③  $f(a) = f(b)$
- ④  $f'(c) = 0 \Rightarrow c \in (a, b)$




## Mean Value Theorem (MVT)

$$f: [a, b] \rightarrow \mathbb{R}$$

- ① Continuous in  $[a, b]$
- ② differentiable in  $(a, b)$
- ③  $f'(c) = \frac{f(b) - f(a)}{b - a}$

$$\Rightarrow c \in (a, b)$$



[Q.1] Verify Rolle's Theorem;  $f(x) = x^2 + 2x - 8$ ,  $x \in [-4, 2]$

$$\left. \begin{array}{l} a = -4 \\ b = 2 \end{array} \right\}$$

we know that

$f(x) = x^2 + 2x - 8$  is continuous in  $[-4, 2]$  and differentiable in  $(-4, 2)$ .

$$f(-4) = (-4)^2 + 2(-4) - 8 = 16 - 8 - 8 = 0$$

$$f(2) = 2^2 + 2(2) - 8 = 8 - 8 = 0$$

$$\therefore \underline{f(-4)} = \underline{f(2)}$$

$$f(x) = x^2 + 2x - 8$$

$$f'(x) = 2x + 2$$

$$f'(c) = 0$$

$$\Rightarrow 2c + 2 = 0$$

$$\Rightarrow \underline{c = -1} \in \underline{(-4, 2)}$$



## Q.2 Rolle's Theorem?

If  $f: [a, b] \rightarrow \mathbb{R}$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$  such that  $f(a) = f(b)$ , then there exists some  $c$  in  $(a, b)$  such that  $f'(c) = 0$ .

## Converse of Rolle's Theorem?

If there exists some  $c$  in  $(a, b)$  such that  $f'(c) = 0$  then  $f$  is continuous on  $[a, b]$  and differentiable on  $(a, b)$  such that  $f(a) = f(b)$ .

Class 11

Maths Reasoning

If  $P$ , then  $Q$   
( $P \rightarrow Q$ )

Converse

If  $Q$  then  $P$   
( $Q \rightarrow P$ )

(i)  $f(x) = [x]$  for  $x \in [5, 9]$

(ii)  $f(x) = [x]$  for  $x \in [-2, 2]$

Similar


(i)  $f(x) = [x]$  for  $x \in [5, 9]$

Slope = 0  
 $\rightarrow 0$

$f(x) = [x]$  is not continuous at integral values of  $x$  in  $[5, 9]$

6, 7, 8, 9

$\therefore$  Rolle's theorem is not applicable.



(Step function)



$f(x) = [x]$  for  $[5, 9]$  Converse of Rolle's Theorem

If  $c \in (a, b), f'(c) = 0$  then

- Continuous  $[a, b]$  ✗
- Diff.  $(a, b)$  ✗
- $f(a) \neq f(b)$  ✗

$c \in (5, 9), f'(c) = 0$

Slope at  $x=c$   
↓  
0

(Slope = 0) → (Graph Horizontal)

$f(5) \neq f(9)$

$[5] \neq [9]$

$5 \neq 9$

Here  $f'(c) = 0$  for some  $c \in (5, 9)$

but  $f$  is not continuous in  $[5, 9]$

∴ Converse of Rolle's theorem is not applicable.

(iii)  $f(x) = x^2 - 1$  for  $x \in [1, 2]$

Rolle's Theorem: Continuous in  $[1, 2]$

Diff. in  $(1, 2)$

$f(a) \neq f(b)$

$f(1) \neq f(2)$   
↓

$1^2 - 1 = 0$

$2^2 - 1 = 3$

Rolle's theorem not applicable



$$f(x) = x^2 - 1 \text{ for } x \in [1, 2]$$

Converse of Rolle's Theorem

If there exists some  $c \in (a, b)$  such that  $f'(c) = 0$  then

- Continuous  $[a, b]$
- Diff.  $(a, b)$
- $f(a) = f(b)$

$$f(x) = x^2 - 1$$

$$f'(x) = 2x$$

$$f'(c) = 2c = 0$$

$$\Rightarrow c = 0 \notin (1, 2)$$

$f'$  does not satisfy the converse of Rolle's Theorem

Q. 3 If  $f: [-5, 5] \rightarrow \mathbb{R}$  is a differentiable function and  $f'(x)$  does not vanish anywhere, ~~where~~  $f(-5) \neq f(5)$ .  
Prove that

Solution

$f$  is a diff. function

then it is continuous.

by MVT

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$c \in (5, 5)$$

$\downarrow$     $\downarrow$   
 $a$     $b$

$$f'(c) = \frac{f(5) - f(-5)}{5 - (-5)} \neq 0$$

$$\Rightarrow f(5) - f(-5) \neq 0$$

$$\Rightarrow f(5) \neq f(-5) \quad \checkmark$$

Vanish = 0

Zero

$$f'(x) \neq 0$$

$$f'(c) \neq 0$$



**Q.4** Verify MVT, if  $f(x) = x^2 - 4x - 3$  in the interval  $[a, b]$ , where  $a=1$  and  $b=4$ .

Ans.  $f(x) = x^2 - 4x - 3$  is continuous in  $[1, 4]$  and differentiable in  $(1, 4)$

Let  $C \in (1, 4)$

$$f'(C) = \frac{f(b) - f(a)}{b - a}$$

$$\Rightarrow 2C - 4 = \frac{(-3) - (-6)}{4 - 1}$$

$$\Rightarrow 2C - 4 = \frac{3}{3} = 1$$

$$\Rightarrow C = \frac{5}{2} \in (1, 4)$$

$$\frac{5}{2} = 2.5$$

$\therefore$  MVT  $\rightarrow$  verified

$$f(x) = x^2 - 4x - 3$$

$$f'(x) = 2x - 4$$

$$f'(C) = 2C - 4$$

$$f(b) = f(4)$$

$$= 4^2 - 4 \cdot 4 - 3$$

$$= -3$$

$$f(a) = f(1)$$

$$= 1^2 - 4 - 3$$

$$= -6$$



Q. 5 Verify MVT, if  $f(x) = x^3 - 5x^2 - 3x$  in the interval  $[a, b]$ , where  $a=1, b=3$ .

Find all  $c \in (1, 3)$  for which  $f'(c) = 0$ .

Sol<sup>n</sup>:  $f$  is continuous in  $[1, 3]$  and differentiable in  $(1, 3)$ .

$a=1, b=3$

$f(x) = x^3 - 5x^2 - 3x$

$f'(x) = 3x^2 - 10x - 3$

$f'(c) = 3c^2 - 10c - 3$

$f(b) = f(3)$

$= 27 - 45 - 9$

$= -29$  (Correct)   
  ~~$-27$  (Wrong)~~

$f(a) = f(1) = 1^3 - 5 - 3 = -7$

$c \in (1, 3)$

$f'(c) = \frac{f(b) - f(a)}{b - a}$

$\Rightarrow 3c^2 - 10c - 3 = \frac{(-29) - (-7)}{3 - 1}$

$\Rightarrow 3c^2 - 10c - 3 = \frac{-22}{2} = -11$

$\Rightarrow 3c^2 - 10c + 8 = 0$

$\Rightarrow 3c^2 - 6c - 4c + 8 = 0$

$\Rightarrow 3c(c-2) - 4(c-2) = 0$

$\Rightarrow (c-2)(3c-4) = 0$

wrong  $c = 2, \frac{4}{3} \in (1, 3)$

Correct  $c = 1, \frac{7}{3} \in (1, 3)$

$f'(c) = 0 \quad c \in (1, 3)$

$\Rightarrow 3c^2 - 10c - 3 = 0 \rightarrow c =$


~~$\Rightarrow 3c^2 - 9c - c - 3 = 0$~~

$\frac{10 \pm \sqrt{100 + 36}}{6} \approx \frac{10 \pm 11.6}{2}$

$\oplus 10.8 \notin (1, 3)$

$\ominus -0.8 \notin (1, 3)$

MVT  
Verified





Verify  
MVT for Q.2 (Parts)

(i)  
 $f(x) = [x], x \in [5, 9]$

not continuous

MVT ✗

(ii)  
 $f(x) = [x]$

$x \in [-2, 2]$

not continuous

MVT ✗

(iii)  $f(x) = x^2 - 1$   
 $x \in [1, 2]$

→ continuous ✓  
[1, 2]

→ Differentiable ✓  
(1, 2)

→  $c \in (1, 2)$

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

$$f(x) = x^2 - 1$$

$$f'(x) = 2x$$

$$f'(c) = 2c$$

$$a = 1, b = 2$$

$$f(a) = f(1) = 0 \quad \checkmark$$

$$f(b) = f(2) = 3 \quad \checkmark$$

$$\Rightarrow 2c = \frac{3 - 0}{2 - 1}$$

$$\Rightarrow 2c = 3$$

$$\Rightarrow \boxed{c = 3/2} \in (1, 2)$$

(1.5) ✓

MVT ✓



Toppers Village Class 12 maths [Continuity and Differentiability]

Miscellaneous Exercise on Chapter-5

**Q.1**  $(3x^2 - 9x + 5)^9$

$\frac{d(x^n)}{dx} = n \cdot x^{n-1}$  (Chain Rule)

$$\frac{d(3x^2 - 9x + 5)^9}{dx} = 9(3x^2 - 9x + 5)^8 \cdot (6x - 9)$$

$$= 27(2x - 3) \cdot (3x^2 - 9x + 5)^8$$

**Q.2**  $y = \sin^3 x + \cos^6 x = (\sin x)^3 + (\cos x)^6$

$$\frac{dy}{dx} = 3(\sin^2 x) \cdot \cos x + 6 \cos^5 x \cdot (-\sin x)$$

$$= 3 \sin x \cos x (\sin x - 2 \cos^4 x)$$

**Q.3**  $y = (5x)^{3 \cos 2x}$

(Let) (Variable) <sup>Variable</sup>

$\Rightarrow \log y = \log (5x)^{3 \cos 2x}$  logarithmic Diff.

$\Rightarrow \log y = 3 \cdot \frac{\cos 2x}{1} \cdot \frac{\log 5x}{1}$   $\frac{d(\log x)}{dx} = \frac{1}{x}$

Diff. w.r.t. (x)  $\rightarrow$

$$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = \left[ -3 \sin 2x \cdot 2 \cdot \log 5x + 3 \cos 2x \cdot \frac{1}{5x} \right]$$

$$\Rightarrow \frac{dy}{dx} = (5x)^{3 \cos 2x} \cdot \left[ -6 \sin 2x \cdot \log 5x + \frac{3 \cos 2x}{x} \right]$$



**Q.4**  $y = \sin^{-1}(x\sqrt{x}) ; 0 \leq x \leq 1$

$$x\sqrt{x} = x^1 \cdot x^{\frac{1}{2}} = x^{\frac{3}{2}}$$

$y = \sin^{-1}(x^{\frac{3}{2}})$   
 diff. w.r.t.  $(x) \rightarrow$  (Chain)  $\frac{d}{dx} x^{(n)} = n \cdot x^{n-1}$

$$\frac{dy}{dx} = \frac{1}{\sqrt{1 - (x^{\frac{3}{2}})^2}} \cdot \frac{3}{2} \cdot x^{\frac{1}{2}}$$

$$\frac{dy}{dx} = \frac{3\sqrt{x}}{2\sqrt{1-x^3}} = \frac{3}{2} \sqrt{\frac{x}{1-x^3}} \quad \checkmark$$

**Q.5**  $y = \frac{\cos^{-1} \frac{x}{2}}{\sqrt{2x+7}}, -2 < x < 2$

$$\left(\frac{u}{v}\right)' = \frac{u'v - u \cdot v'}{v^2}$$

$$(u \cdot v)' = u' \cdot v + u \cdot v'$$

$$\Rightarrow y = \cos^{-1} \frac{x}{2} \cdot \frac{1}{\sqrt{2x+7}}$$

$$\Rightarrow y = \underbrace{\cos^{-1} \frac{x}{2}} \cdot \underbrace{(2x+7)^{-\frac{1}{2}}}$$

$$\Rightarrow \frac{dy}{dx} = \underbrace{-\frac{1}{\sqrt{1 - \left(\frac{x}{2}\right)^2}} \cdot \frac{1}{2}} \cdot (2x+7)^{-\frac{1}{2}} + \cos^{-1} \frac{x}{2} \cdot \underbrace{\left[-\frac{1}{2} (2x+7)^{-\frac{1}{2}-1} \cdot 2\right]}$$

$$\Rightarrow \frac{dy}{dx} = -\frac{1}{\sqrt{\frac{4-x^2}{4}} \cdot \sqrt{2x+7}} - \frac{\cos^{-1}(\frac{x}{2})}{(2x+7)^{3/2}}$$

$$\frac{dy}{dx} = -\left[ \frac{1}{\sqrt{4-x^2} \cdot \sqrt{2x+7}} + \frac{\cos^{-1} \frac{x}{2}}{(2x+7)^{3/2}} \right] \quad \checkmark$$



Q.6.

$$y = \cot^{-1} \left[ \frac{\sqrt{1+\sin x} + \sqrt{1-\sin x}}{\sqrt{1+\sin x} - \sqrt{1-\sin x}} \right], \quad 0 < x < \frac{\pi}{2}$$

$$\checkmark \quad 1 + \sin x = \sin^2 \frac{x}{2} + \cos^2 \frac{x}{2} + 2 \sin \frac{x}{2} \cdot \cos \frac{x}{2} = \left( \sin \frac{x}{2} + \cos \frac{x}{2} \right)^2$$

$$\checkmark \quad 1 - \sin x = \sin^2 \frac{x}{2} + \cos^2 \frac{x}{2} - 2 \sin \frac{x}{2} \cdot \cos \frac{x}{2} = \left( \sin \frac{x}{2} - \cos \frac{x}{2} \right)^2$$

$$\checkmark \quad \sqrt{x^2} = |x|$$

$$(\sqrt{x})^2 = x$$

$$|x| = \begin{cases} x, & x \geq 0 \quad (+) \\ -x, & x < 0 \quad (-) \end{cases}$$


$$y = \cot^{-1} \left( \frac{\sqrt{1+\sin x} + \sqrt{1-\sin x}}{\sqrt{1+\sin x} - \sqrt{1-\sin x}} \right), \quad 0 < x < \frac{\pi}{2}$$

$$y = \cot^{-1} \left( \frac{\sqrt{\left( \sin \frac{x}{2} + \cos \frac{x}{2} \right)^2} + \sqrt{\left( \sin \frac{x}{2} - \cos \frac{x}{2} \right)^2}}{\sqrt{\left( \sin \frac{x}{2} + \cos \frac{x}{2} \right)^2} - \sqrt{\left( \sin \frac{x}{2} - \cos \frac{x}{2} \right)^2}} \right)$$

$$y = \cot^{-1} \left( \frac{\left| \sin \frac{x}{2} + \cos \frac{x}{2} \right| + \left| \sin \frac{x}{2} - \cos \frac{x}{2} \right|}{\left| \sin \frac{x}{2} + \cos \frac{x}{2} \right| - \left| \sin \frac{x}{2} - \cos \frac{x}{2} \right|} \right)$$

$$0 < x < \frac{\pi}{2}$$

$$0 < \frac{x}{2} < \frac{\pi}{4}$$



$$\cos \frac{x}{2} > \sin \frac{x}{2}$$

$$\star \quad \left| \sin \frac{x}{2} - \cos \frac{x}{2} \right| = -(\sin \frac{x}{2} - \cos \frac{x}{2})$$



$$\Rightarrow y = \cot^{-1} \left( \frac{(\cancel{\sin \frac{x}{2}} + \cos \frac{x}{2}) - (\cancel{\sin \frac{x}{2}} - \cos \frac{x}{2})}{(\cancel{\sin \frac{x}{2}} + \cos \frac{x}{2}) + (\cancel{\sin \frac{x}{2}} - \cos \frac{x}{2})} \right)$$

$$\Rightarrow y = \cot^{-1} \left( \frac{\cancel{2} \cos \frac{x}{2}}{\cancel{2} \sin \frac{x}{2}} \right)$$

$$\Rightarrow y = \cot^{-1} \left( \cot \frac{x}{2} \right)$$

$$\Rightarrow y = \frac{x}{2}$$

Diff. w.r.t. (x)  $\rightarrow$

$$\frac{dy}{dx} = \frac{1}{2}$$



Miscellaneous Exercise on Chapter-5

[Q.7]  $y = (\log x)^{\log x}, x > 1$

$$\Rightarrow \log y = \log (\log x)^{\log x}$$

$$\Rightarrow \log y = \log x \cdot \log (\log x)$$

by Diff. w.r.t.  $x \rightarrow$

$$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = \left[ \frac{1}{x} \cdot \log (\log x) + \log x \cdot \left( \frac{1}{\log x} \cdot \frac{1}{x} \right) \right]$$

$$\Rightarrow \frac{dy}{dx} = (\log x)^{\log x} \cdot \left[ \frac{\log (\log x)}{x} + \frac{1}{x} \right]$$

logarithmic  
Differentiation

V.V.

log, Diff

$$\frac{d(\log x)}{dx} = \frac{1}{x}$$

[Q.8]  $y = \cos(a \cos x + b \sin x)$   
(Chain Rule)

$a, b \rightarrow \text{constants.}$

$$\frac{dy}{dx} = -\sin(a \cos x + b \sin x) \cdot (-a \sin x + b \cos x)$$

$$= (a \sin x - b \cos x) \cdot \sin(a \cos x + b \sin x)$$



$$\boxed{Q.9} \quad y = (\sin x - \cos x)^{(\sin x - \cos x)}, \quad \frac{\pi}{4} < x < \frac{3\pi}{4}$$

$$\Rightarrow \log y = \log \left[ \frac{(\sin x - \cos x)}{(\sin x - \cos x)} \right] \quad (\log m^n = n \cdot \log m)$$

$$\Rightarrow \log y = (\sin x - \cos x) \cdot \log (\sin x - \cos x)$$

diff. w.r.t.  $x \rightarrow$

$$\Rightarrow \frac{1}{y} \cdot \frac{dy}{dx} = \left[ \underbrace{(\cos x + \sin x)}_{\checkmark} \cdot \log (\sin x - \cos x) + \underbrace{(\sin x - \cos x)}_{\cancel{\sin x - \cos x}} \cdot \left( \frac{1}{\cancel{(\sin x - \cos x)}} \cdot \underbrace{(\cos x + \sin x)}_{\checkmark} \right) \right]$$

$$\Rightarrow \frac{dy}{dx} = (\sin x - \cos x)^{(\sin x - \cos x)} \cdot (\cos x + \sin x)$$

$$\left[ \log (\sin x - \cos x) + 1 \right]$$



**Q.10**  $y = x^n + x^a + a^x + a^a$

(Fixed)  
 $a > 0$   
 $x > 0$

$$\frac{dy}{dx} = \frac{d(x^n)}{dx} + \frac{d(x^a)}{dx} + \frac{d(a^x)}{dx} + \frac{d(a^a)}{dx}$$

$$\frac{dy}{dx} = \frac{d(x^n)}{dx} + a \cdot x^{a-1} + \frac{d(a^x)}{dx} + 0 \quad \text{--- (1) } a^a = \text{const.}$$

$$x^a \rightarrow (x^n) \rightarrow n \cdot x^{n-1}$$

$$\frac{d(a^x)}{dx} = a^x \cdot \log a$$

$u = x^n$  (let)

$\Rightarrow \log u = \log x^n$

$\Rightarrow \log u = \underline{x \cdot \log x}$

diff. w.r.t.  $(x) \rightarrow$

$\Rightarrow \frac{1}{u} \cdot \frac{du}{dx} = \left( 1 \cdot \log x + \cancel{x} \cdot \frac{1}{\cancel{x}} \right)$

$\Rightarrow \frac{du}{dx} = x^n (\log x + 1)$

$\Rightarrow \frac{d(x^n)}{dx} = \underline{x^n (\log x + 1)}$

$v = a^x$

$\Rightarrow \log v = x \cdot \log a$

$\Rightarrow \frac{dv}{v} = 1 \cdot \log a$

$\Rightarrow \frac{dv}{dx} = a^x \cdot \log a$

$\Rightarrow \frac{dv}{dx} = a^x \cdot \log a$

By eq<sup>n</sup> (1)  $\rightarrow$

$$\frac{dy}{dx} = \underline{x^n (\log x + 1)} + \underline{a \cdot x^{a-1}} + \underline{a^x \cdot \log a} + \underline{0}$$



**Q.11**  $y = x^{x^2-3} + (x-3)^{x^2}$  , for  $x > 3$

$$\boxed{y = u + v} \Rightarrow \boxed{\frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}}$$

$$u = x^{x^2-3}$$

$$\Rightarrow \log u = \log (x^{x^2-3})$$

$$\Rightarrow \log u = (x^2-3) \cdot \log x$$

(diff. w.r.t.  $x$ )  $\rightarrow$

$$\Rightarrow \frac{1}{u} \cdot \left( \frac{du}{dx} \right) = \left[ \underline{(2x)} \cdot \log x + (x^2-3) \cdot \underline{\left( \frac{1}{x} \right)} \right]$$

$$\Rightarrow \frac{du}{dx} = x^{x^2-3} \cdot \left[ 2x \log x + \frac{x^2-3}{x} \right]$$

$$\therefore \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$$

$$\Rightarrow \frac{dy}{dx} = x^{x^2-3} \cdot \left[ 2x \cdot \log x + \frac{x^2-3}{x} \right] + (x-3)^{x^2} \cdot \left[ 2x \cdot \log(x-3) + \frac{x^2}{x-3} \right]$$

$$v = (x-3)^{x^2}$$

$$\Rightarrow \log v = \log (x-3)^{x^2}$$

$$\Rightarrow \log v = x^2 \cdot \log(x-3)$$

Diff. w.r.t.  $x$   $\rightarrow$

$$\Rightarrow \frac{1}{v} \cdot \frac{dv}{dx} = \left[ \underline{(2x)} \cdot \log(x-3) + x^2 \cdot \underline{\frac{1}{x-3}} \right]$$

$$\Rightarrow \frac{dv}{dx} = (x-3)^{x^2} \cdot \left[ 2x \cdot \log(x-3) + \frac{x^2}{x-3} \right]$$



Miscellaneous Exercise on Chapter 5

Q.12 Find  $\frac{dy}{dx}$ , if  $y = 12(1 - \cos t)$ ,  $x = 10(t - \sin t)$ ;   
 (Parametric Form)  $-\frac{\pi}{2} < t < \frac{\pi}{2}$

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)} = \frac{\left(\frac{d(12(1 - \cos t))}{dt}\right)}{\left(\frac{d(10(t - \sin t))}{dt}\right)}$$

$$= \frac{12(0 + \sin t)}{10(1 - \cos t)} = \frac{6(\sin t)}{5(1 - \cos t)}$$

$$\left( \begin{array}{l} \sin t = 2 \sin t_{/2} \cdot \cos t_{/2} \\ 1 - \cos t = 2 \sin^2 t_{/2} \end{array} \right) \rightarrow = \frac{6 \cdot \cancel{2} \sin t_{/2} \cdot \cos t_{/2}}{5 (\cancel{2} \sin^2 t_{/2})}$$

$$\frac{dy}{dx} = \frac{6}{5} \cot\left(\frac{t}{2}\right)$$

Q.13 Find  $\frac{dy}{dx}$ , if  $y = \sin^{-1} x + \sin^{-1}(\sqrt{1-x^2})$ ,  $0 < x < 1$

I-method.

$$y = \sin^{-1} x + \sin^{-1}(\sqrt{1-x^2}) \quad \text{(Chain Rule)}$$

$$\frac{dy}{dx} = \frac{1}{\sqrt{1-x^2}} + \frac{1}{\sqrt{1-(\sqrt{1-x^2})^2}} \times \frac{1}{\cancel{\sqrt{1-x^2}}} (0 - \cancel{x})$$



$$\Rightarrow \frac{dy}{dx} = \frac{1}{\sqrt{1-x^2}} - \frac{x}{\sqrt{1-(1-x^2)} \cdot \sqrt{1-x^2}}$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{\sqrt{1-x^2}} - \frac{x}{\sqrt{x-x+x^2} \cdot \sqrt{1-x^2}}$$

$$\Rightarrow \frac{dy}{dx} = + \frac{1}{\sqrt{1-x^2}} - \frac{x}{x \cdot \sqrt{1-x^2}}$$

$$\frac{dy}{dx} = 0 \quad \checkmark$$

II - method

$$y = \sin^{-1}(x) + \sin^{-1} \sqrt{1-x^2}$$

Inverse trigonometry

Substitution,  $x = \sin \theta$

$$y = \sin^{-1}(\sin \theta) + \sin^{-1} \sqrt{1 - \sin^2 \theta}$$

$$\Rightarrow y = \theta + \sin^{-1} \sqrt{\cos^2 \theta}$$

$$\Rightarrow y = \theta + \sin^{-1}(\cos \theta)$$

$$\sin(90^\circ - \theta) = \cos \theta$$

$$\Rightarrow y = \theta + \sin^{-1}(\sin(\frac{\pi}{2} - \theta))$$

$$\Rightarrow y = \cancel{\theta} + \frac{\pi}{2} - \cancel{\theta}$$

$$\Rightarrow y = \frac{\pi}{2} \leftarrow \text{Constant}$$

$$\boxed{\frac{dy}{dx} = 0}$$



**Q.14**

If  $x\sqrt{1+y} + y\sqrt{1+x} = 0$ , for  $-1 < x < 1$ ,

Prove that  $\frac{dy}{dx} = -\frac{1}{(1+x)^2}$

Ans.

$$x\sqrt{1+y} + y\sqrt{1+x} = 0$$

$$\Rightarrow x\sqrt{1+y} = -y\sqrt{1+x}$$

Square both sides.

$$\Rightarrow x^2(1+y) = y^2(1+x)$$

$$\Rightarrow x^2 + x^2y = y^2 + xy^2$$

$$\Rightarrow (x^2 - y^2) + (x^2y - xy^2) = 0$$

$$\Rightarrow (x+y)(x-y) + xy(x-y) = 0$$

$$\Rightarrow (x-y) \{ \underline{x+y + xy} \} = 0$$

$x=y$ ,  $x+y+xy=0$

$y=x$

I

$$\Rightarrow y(1+x) = -x$$

$$\Rightarrow y = \frac{-x}{1+x}$$

II

original eqn. Cross check

$$x\sqrt{1+y} + y\sqrt{1+x} = 0$$

I  $y=x$

$$\Rightarrow x\sqrt{1+x} + x\sqrt{1+x} = 0$$

$$\Rightarrow 2x\sqrt{1+x} = 0 \Rightarrow x=0, x=-1$$

Constants

Reject

original (Equation)<sup>2</sup>  
↓  
Extra roots.  
Satisfy that

$y=x \rightarrow 0$   
 $y=x \rightarrow -1$   
 $y=0$   
 $y=-1$



Now we will check  $y = \frac{-x}{1+x}$  in the

original equation  $\rightarrow x\sqrt{1+y} + y\sqrt{1+x} = 0$

$$\Rightarrow x\sqrt{1 - \frac{x}{1+x}} - \frac{x}{1+x}\sqrt{1+x} = 0$$

$$\Rightarrow x\sqrt{\frac{1+x-x}{1+x}} - \frac{x}{\sqrt{1+x}} = 0$$

$$\Rightarrow x\sqrt{\frac{1}{1+x}} - \frac{x}{\sqrt{1+x}} = 0$$

$$\Rightarrow \frac{x}{\sqrt{1+x}} - \frac{x}{\sqrt{1+x}} = 0$$

$$\Rightarrow 0 = 0$$

$$y = \frac{-x}{1+x}$$

Accepted

by Diff. w.r.t.  $x \rightarrow$

$$\left(\frac{u}{v}\right)' = \frac{u'v + u.v'}{v^2}$$

$$\frac{dy}{dx} = \frac{(-1) \cdot (1+x) - (-x) \cdot (1)}{(1+x)^2}$$

$$\frac{dy}{dx} = \frac{-1-x+x}{(1+x)^2} = \frac{-1}{(1+x)^2}$$



Miscellaneous Exercise on Chapter (5)

[Q.15] If  $(x-a)^2 + (y-b)^2 = c^2$ , for some  $c > 0$ ,

Prove that  $\frac{\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{3/2}}{\left(\frac{d^2y}{dx^2}\right)}$  is a constant independent of 'a' and 'b'.

Sol<sup>n</sup>  $(x-a)^2 + (y-b)^2 = c^2$

by diff. w.r.t. 'x' →

a, b, c → Constant

$$\Rightarrow 2(x-a) + 2(y-b) \cdot \frac{dy}{dx} = 0$$

$$\Rightarrow (x-a) + (y-b) \cdot \frac{dy}{dx} = 0$$

by diff. w.r.t. 'x' →

$$\frac{dy}{dx} = \frac{-(x-a)}{(y-b)}$$

$$\Rightarrow 1 + \left(\frac{dy}{dx} - 0\right) \cdot \frac{dy}{dx} + (y-b) \cdot \frac{d^2y}{dx^2} = 0$$

$$\Rightarrow 1 + \left(\frac{dy}{dx}\right)^2 + (y-b) \cdot \frac{d^2y}{dx^2} = 0$$

$$\Rightarrow (y-b) \cdot \left(\frac{d^2y}{dx^2}\right) = - \left\{1 + \left(\frac{dy}{dx}\right)^2\right\}$$

$$\Rightarrow \frac{d^2y}{dx^2} = \frac{- \left\{1 + \left(\frac{dy}{dx}\right)^2\right\}}{y-b}$$



$$1 + \left(\frac{dy}{dx}\right)^2 = 1 + \left[-\left(\frac{x-a}{y-b}\right)\right]^2 = 1 + \frac{(x-a)^2}{(y-b)^2}$$

$$= \frac{(y-b)^2 + (x-a)^2}{(y-b)^2} = \frac{c^2}{(y-b)^2}$$

Given:  $(x-a)^2 + (y-b)^2 = c^2$

$$\Rightarrow \boxed{1 + \left(\frac{dy}{dx}\right)^2 = \frac{c^2}{(y-b)^2}}$$

$$\frac{d^2y}{dx^2} = - \frac{\left\{1 + \left(\frac{dy}{dx}\right)^2\right\}}{(y-b)} = - \frac{\left\{\frac{c^2}{(y-b)^2}\right\}}{(y-b)}$$

$$\boxed{\frac{d^2y}{dx^2} = - \frac{c^2}{(y-b)^3}}$$

$$\frac{\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{3/2}}{\frac{d^2y}{dx^2}} = \frac{\left[\frac{c^2}{(y-b)^2}\right]^{3/2}}{- \frac{c^2}{(y-b)^3}} = \frac{\cancel{c^2}^c}{\cancel{(y-b)^3}^3} = -c$$

Constant, independent of  $a, b$



**Q.16** If  $\cos y = (x) \cos(a+y)$ , with  $\cos a \neq \pm 1$ ,

Prove that  $\frac{dy}{dx} = \frac{\cos^2(a+y)}{\sin a}$ .

Ans.  $\cos y = x \cdot \cos(a+y)$   $(u \cdot v)' = u' \cdot v + u \cdot v'$   
diff. w.r.t.  $(x) \rightarrow$

$$\Rightarrow -\sin y \cdot \left(\frac{dy}{dx}\right) = 1 \cdot \cos(a+y) - x \sin(a+y) \cdot \left(\frac{dy}{dx}\right)$$

$$\Rightarrow \left(\frac{dy}{dx}\right) [x \cdot \sin(a+y) - \sin y] = \cos(a+y)$$

$$\Rightarrow \frac{dy}{dx} = \frac{\cos(a+y)}{(x) \cdot \sin(a+y) - \sin y}$$

Given  $\cos y = (x) \cdot \cos(a+y)$

$$x = \frac{\cos y}{\cos(a+y)}$$

$$\Rightarrow \frac{dy}{dx} = \frac{\cos(a+y)}{\frac{\cos y}{\cos(a+y)} \cdot \sin(a+y) - \sin y}$$

$$\Rightarrow \frac{dy}{dx} = \frac{\cos(a+y)}{\cos y \cdot \sin(a+y) - \sin y \cdot \cos(a+y)}$$

$$\frac{\sin A \cdot \cos B - \sin B \cdot \cos A}{\downarrow \quad \downarrow} = \sin(A-B) = \sin(a+y-y)$$



$$\frac{dy}{dx} = \frac{\cos^2(a+y)}{\sin[(a+y)-y]} \leftarrow \text{By } \sin(A-B)$$

$$\frac{dy}{dx} = \frac{\cos^2(a+y)}{\sin a}$$

Hence Proved.

**Q.17** If  $x = a(\cos t + t \sin t)$  and

$y = a(\sin t - t \cos t)$ , find  $\frac{d^2y}{dx^2}$ .

$$\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)}$$

$$\frac{dy}{dx} = \frac{a t \sin t}{a t \cos t} = \tan t$$

$$\boxed{\frac{dy}{dx} = \tan t}$$

diff. w.r.t. 'x' (again)

$$\Rightarrow \frac{d^2y}{dx^2} = \sec^2 t \cdot \left(\frac{dt}{dx}\right)$$

$$\Rightarrow \frac{d^2y}{dx^2} = \sec^2 t \cdot \frac{1}{a t \cos t}$$

$$\boxed{\frac{d^2y}{dx^2} = \frac{\sec^3 t}{a t}}$$

$$\frac{dy}{dt} = a(\cancel{\cos t} - 1 \cdot \cancel{\cos t} + t \cdot \sin t)$$

$$\frac{dy}{dt} = a t \sin t$$

$$\frac{dx}{dt} = a(-\cancel{\sin t} + 1 \cdot \cancel{\sin t} + t \cdot \cos t)$$

$$\frac{dx}{dt} = a t \cos t$$

Reciprocal,

$$\frac{dt}{dx} = \frac{1}{a t \cos t}$$



Miscellaneous Exercise on chapter 5

**Q.18** If  $f(x) = |x|^3$ , show that  $f''(x)$  exists for all real  $x$  and find it.

Sol<sup>n</sup>  $|x|$  changes its nature at  $x=0$  

$f$  differentiable  $\Rightarrow f'$  exists

$f'$  differentiable  $\Rightarrow f''$  exists

Continuity  
yes

LHD = RHD

$$f'(0^-) = f'(0^+)$$

LHD = RHD

$$f''(0^-) = f''(0^+)$$

$$f(x) = |x|^3$$

$$f(x) = \begin{cases} x^3, & x \geq 0 \\ -x^3, & x < 0 \end{cases}$$

$$f'(x) = \begin{cases} 3x^2, & x \geq 0 \rightarrow \text{RHD of } f = f'(0^+) = 3(0)^2 = 0 \\ -3x^2, & x < 0 \rightarrow \text{LHD of } f = f'(0^-) = -3(0)^2 = 0 \end{cases}$$

$$f''(x) = \begin{cases} 6x, & x \geq 0 \rightarrow \text{RHD of } f' = f''(0^+) = 6(0) = 0 \\ -6x, & x < 0 \rightarrow \text{LHD of } f' = f''(0^-) = -6(0) = 0 \end{cases}$$

$$f''(x) = 6x$$

$f'' \rightarrow \text{exists}$



Q.19

Prove by PMI

$$\frac{d(x^n)}{dx} = n \cdot x^{n-1}$$

for all  
positive  
integers  $n$ .

$P(n)$ : Statement

①  $P(1)$  true (Prove)

② let  $P(k)$  is true  
then prove  $P(k+1)$  is also true

$$n \in \{1, 2, 3, \dots\}$$

$$P(n): \frac{d(x^n)}{dx} = n \cdot x^{n-1}$$

Step-① we have to prove that  $P(1)$  is true.

$$P(1): \frac{d(x^1)}{dx} = 1 \cdot x^{1-1}$$

$$1 = 1 \cdot x^0$$

$$1 = 1 \quad \text{(true)}$$

Step-② Let  $P(k)$  is true

$$P(k): \frac{d(x^k)}{dx} = k \cdot x^{k-1} \quad \text{--- (1)}$$

Now We have to prove that  $P(k+1)$  is true.

i.e.

$$P(k+1): \frac{d(x^{k+1})}{dx} = (k+1) \cdot x^k$$



$$\text{LHS of } P(k+1) = \frac{d(x^{k+1})}{dx} = \frac{d(\underbrace{x^k}_{u} \cdot \underbrace{x}_v)}{dx}$$

$$= \left( \frac{d(x^k)}{dx} \right) \cdot x + x^k \cdot \frac{d(x)}{dx}$$

↓ exp

$$= (k \cdot \underline{x^{k-1}}) \cdot \underline{x'} + x^k \cdot (1)$$

$$= k \cdot \underline{x^k} + \underline{x^k}$$

$$= (k+1) x^k = \text{RHS of } P(k+1)$$

$\therefore P(k+1)$  is also true.

$$\Rightarrow P(n): \underbrace{\frac{d(x^n)}{dx} = n \cdot x^{n-1}}_{\text{is true.}} \quad \underline{\underline{n \in \mathbb{N}}}$$

[Q.20] Using the fact that  $\sin(A+B) = \sin A \cos B + \cos A \sin B$  and the differentiation, obtain the sum formula for cosines.  $\rightarrow$   $\cos(A+B) = ?$

Ans.

$$\sin(A+B) = \sin A \cdot \cos B + \cos A \cdot \sin B$$

A & B  $\rightarrow$  variable

$\rightarrow$  by Differentiating with respect to 'x'  $\rightarrow$

$$\Rightarrow \underline{\underline{\cos(A+B)}} \cdot \left( \frac{dA}{dx} + \frac{dB}{dx} \right) = \cos A \cdot \frac{dA}{dx} \cdot \cos B + \sin A \cdot \sin B \cdot \frac{dB}{dx} - \sin A \cdot \frac{dA}{dx} \cdot \sin B + \cos A \cdot \cos B \cdot \frac{dB}{dx}$$



$$\Rightarrow \cos(A+B) \cdot \left( \frac{dA}{dn} + \frac{dB}{dn} \right) = \cos A \cdot \frac{dA}{dn} \cdot \cos B - \sin A \sin B \cdot \frac{dB}{dn} \\ - \sin A \cdot \frac{dA}{dn} \cdot \sin B + \cos A \cdot \cos B \cdot \frac{dB}{dn}$$

$$\Rightarrow \frac{dA}{dn} (\cos A \cos B - \sin A \sin B) \\ + \frac{dB}{dn} (\cos A \cos B - \sin A \sin B)$$

$$\Rightarrow \cos(A+B) = \left( \frac{dA}{dn} + \frac{dB}{dn} \right) (\cos A \cos B - \sin A \sin B)$$

$$\Rightarrow \boxed{\cos(A+B) = \cos A \cos B - \sin A \sin B}$$

**[Q.2]** Does there exist a function which is continuous everywhere but not differentiable at exactly two points? Justify.

Ans.






Graphically

Differentiable

Non Differentiable

Graph Smooth


Sharp corners





Continuous everywhere

(Non differentiable  
at exactly 2 points.)



(exactly 2 sharp corner)

$$y = mx + c$$

↑  
(slope)

$$y = -x \quad x \in (-\infty, 0)$$

$$y = x \quad x \in [0, 1]$$

$$y = 1 \quad x \in (1, \infty)$$

$$f(x) = \begin{cases} -x, & x < 0 \\ x, & 0 \leq x \leq 1 \\ 1, & x > 1 \end{cases}$$

Continuous everywhere

but not differentiable at exactly  
2 points.

$$x = 0, 1$$



Toppers Village Class 12 maths [Continuity and Differentiability]

Miscellaneous Exercise on Chapter 5

Q.22. If  $y = \begin{vmatrix} f(n) & g(n) & h(n) \\ l & m & n \\ a & b & c \end{vmatrix}$ , prove that

$$\frac{dy}{dn} = \begin{vmatrix} f'(n) & g'(n) & h'(n) \\ l & m & n \\ a & b & c \end{vmatrix}$$

$f(n), g(n), h(n)$   
variable  
 $(l, m, n)$   
 $(a, b, c) \rightarrow \text{constant}$

Solution.

$$y = \begin{vmatrix} f(n) & g(n) & h(n) \\ l & m & n \\ a & b & c \end{vmatrix} \rightarrow R_1 \text{ as along expand}$$

$$y = \underbrace{f(n)}_{\text{Const.}} \begin{vmatrix} m & n \\ b & c \end{vmatrix} - \underbrace{g(n)}_{\text{Const.}} \begin{vmatrix} l & n \\ a & c \end{vmatrix} + \underbrace{h(n)}_{\text{Const.}} \begin{vmatrix} l & m \\ a & b \end{vmatrix}$$

diff. w.r.t.  $(n) \rightarrow$

$$\frac{dy}{dn} = \underbrace{f'(n)}_{\text{Const.}} \begin{vmatrix} m & n \\ b & c \end{vmatrix} - \underbrace{g'(n)}_{\text{Const.}} \begin{vmatrix} l & n \\ a & c \end{vmatrix} + \underbrace{h'(n)}_{\text{Const.}} \begin{vmatrix} l & m \\ a & b \end{vmatrix}$$

$$\frac{dy}{dn} = \begin{vmatrix} f'(n) & g'(n) & h'(n) \\ l & m & n \\ a & b & c \end{vmatrix}$$

✓ ✓ ✓ ~~1~~ ~~2~~ ~~3~~



Note :

$$\begin{vmatrix} f_1 & f_2 & f_3 \\ g_1 & g_2 & g_3 \\ h_1 & h_2 & h_3 \end{vmatrix}' = \begin{vmatrix} f_1' & f_2' & f_3' \\ g_1 & g_2 & g_3 \\ h_1 & h_2 & h_3 \end{vmatrix} + \begin{vmatrix} f_1 & f_2 & f_3 \\ g_1' & g_2' & g_3' \\ h_1 & h_2 & h_3 \end{vmatrix} + \begin{vmatrix} f_1 & f_2 & f_3 \\ g_1 & g_2 & g_3 \\ h_1' & h_2' & h_3' \end{vmatrix}$$

information for Q.22

[Q.23] If  $y = e^{a \cos^{-1} x}$ ,  $-1 \leq x \leq 1$ , show that

Ans.  $y = e^{a \cos^{-1} x}$

Diff. w.r.t.  $(x) \rightarrow$

Chain Rule

$$\Rightarrow \frac{dy}{dx} = e^{a \cos^{-1} x} \cdot a \left( \frac{-1}{\sqrt{1-x^2}} \right)$$

$$\Rightarrow \sqrt{1-x^2} \cdot \frac{dy}{dx} = -a y$$

by squaring  $\rightarrow$

$$\Rightarrow (1-x^2) \cdot \left( \frac{dy}{dx} \right)^2 = a^2 y^2$$

by diff. w.r.t.  $(x) \rightarrow$

$$\Rightarrow (-2x) \cdot \left( \frac{dy}{dx} \right)^2 + (1-x^2) \cdot 2 \left( \frac{dy}{dx} \right) \cdot \frac{d^2y}{dx^2} = a^2 \cdot 2y \cdot \frac{dy}{dx}$$

$$(1-x^2) \cdot \frac{d^2y}{dx^2} - x \frac{dy}{dx} - a^2 y = 0$$

$$(1-x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} - a^2 y = 0$$

Hence proved



## **All Useful Links:**

- **YouTube Channel : Toppers Village (Get full and free video lectures of Class 10 & 11 & 12 Maths) -**  
<https://www.youtube.com/toppersvillage>
- **Website :** [www.toppersvillage.com](http://www.toppersvillage.com)
- **Instagram :**  
<https://www.instagram.com/toppersvillage/>
- **Facebook Page:** [www.facebook.com/ToppersVillage/](https://www.facebook.com/ToppersVillage/)
- **Facebook Group:**  
<https://www.facebook.com/groups/ToppersVillage>